

N/C

00700642  
AUG 2000 (21.08.00)



Europäisches  
Patentamt

Eur pean  
Patent Office

Office eur péen  
des brevets

CA00642

REC'D 13 SEP 2000	
WIPO	PCT

4

Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

99201745.9

**PRIORITY DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH  
RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts:  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

J.L.C. HATTEN-HECKMAN

DEN HAAG, DEN  
THE HAGUE, 20/06/00  
LA HAYE, LE

10 60 70 114



Europäisches  
Patentamt

European  
Patent Office

Office eur. p<sup>at</sup>ent  
des brevets

**Blatt 2 der Bescheinigung**  
**Sheet 2 of the certificate**  
**Page 2 de l'attestation**

Anmeldung Nr.: 99201745.9  
Application no.:  
Demande n°:

Anmeldetag:  
Date of filing:  
Date de dépôt: 02/06/99

Anmelder:  
Applicant(s):  
Demandeur(s):  
CENTRUM VOOR PLANTENVEREDELINGS- EN REPRODUCTIEONDERZOEK  
6708 PB Wageningen  
NETHERLANDS

Her Majesty in Right of Canada, represented by The Minister of Agriculture and Agri-Food Canada  
Ottawa, Ontario K1A 0G6

CANADA  
Bezeichnung der Erfindung:  
Title of the invention:  
Titre de l'invention:

Use of the BNM3 transcriptional activator to control plant embryogenesis and regeneration processes

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat:  
State:  
Pays:

Tag:  
Date:  
Date:


Aktenzeichen:  
File no.  
Numéro de dépôt:

Internationale Patentklassifikation:  
International Patent classification:  
Classification internationale des brevets:

C12N15/29, C12N15/82, C12N5/10, C07K14/415, A01H5/00, A01H5/10

Am Anmeldetag benannte Vertragsstaaten:  
Contracting states designated at date of filing: AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GRIE/IT/LU/LU/MC/NL/PT/SE  
Etats contractants désignés lors du dépôt:

Bemerkungen:  
Remarks:  
Remarques:



- 1 -

EPO - DG 1

USE OF THE *BNM3* TRANSCRIPTIONAL ACTIVATOR TO CLONAL PLANT  
EMBRYOGENESIS AND REGENERATION PROCESSES

(58)

The present invention relates to asexual embryo formation and regeneration in plants. More specifically, it relates to processes for producing asexually-derived embryos, and for enhancing regeneration capacity in plants. The present invention also relates to heterologous protein production systems in plants, and the uses thereof.

## BACKGROUND OF THE INVENTION

A typical angiosperm seed consists of three major components, the embryo, the endosperm and the maternal seed coat. Seed development begins with a double fertilization event, in which one sperm cell nucleus fuses with the egg cell nucleus to form the embryo, and a second sperm cell nucleus fuses with two central cell nuclei to form the endosperm. Embryo development itself can be separated into three developmental phases. The first phase of embryo development is one of cell division and morphogenesis, which serves to establish the major tissue types and organ systems of the mature plant. The second phase encompasses a period of rapid cell expansion and is characterized by the synthesis of storage reserves that sustain the embryo during germination and early seedling development. In the final phase of embryo development, the embryo becomes desiccated and enters into a period of developmental arrest or dormancy. ~~All of the above events normally take place while the seed remains attached to the maternal plant.~~

Many plant species are capable of producing embryos in the absence of fertilization. This process of asexual embryo development may occur naturally, for example on the leaf margins of *Bryophyllum* (Yarborough, 1923) and *Malaxis* (Taylor, 1967), or within the ovule of apomictic plants (Koltunow, 1995). Apomixis refers to the production of a seed from the maternal ovule tissues in the absence of egg cell fertilization. Asexual embryo development may also be induced *in vitro* from gametophytic or somatic tissue (Mordhorst *et al.*, 1997) or, as shown recently, may be

induced by genetic modification of gene expression (Ogas *et al.*, 1997; Lotan *et al.*, 1998).

Three major mechanisms of apomixis, diplospory, apospory and adventitious embryony, have been observed. Each mechanism differs with respect to the source of the cell that gives rise to the embryo and with respect to the time during ovule development at which the apomictic process is initiated. Diplospory and apospory are considered gametophytic forms of apomixis as they involve the formation of diploid embryo sacs. Adventitious embryony does not involve the production of a mitotically-derived embryo sac.

In diplospory, the megaspore mother cell does not undergo normal meiosis, but rather divides mitotically to produce a diploid embryo sac instead of the normal haploid embryo sac. One of the cells of the embryo sac functions as the egg cell and divides parthenogenetically (without fertilization) to form an embryo. In some species the unreduced polar nuclei of the embryo sac may fuse to form the endosperm (autonomous endosperm production), the nutritive tissue of the seed, while in other species pollination is necessary for endosperm production (pseudogamy).

In aposporous apomicts, parthenogenic embryos are produced from additional cells, the aposporous initials, that differentiate from the nucellus. As with the megagametophyte of diplosporous species, the aposporous initial undergoes mitotic divisions to produce a diploid embryo sac. Aposporous embryos are not derived from the megagametophyte and can therefore co-exist within a single ovule with sexually-derived embryos. Autonomous production of endosperm is rare in aposporous species. Aposporous apomicts therefore depend on fertilization of the polar nuclei of a meiotically-derived embryo sac for the production of endosperm.

With adventitious embryony, embryos are formed directly from sporophytic ovule tissue, such as the integuments or nucellus, via parthenogenesis. Seeds derived from species exhibiting adventitious embryony generally contain multiple asexually-derived

embryos and may also contain a single sexually-derived embryo. Plants exhibiting adventitious embryo also rely on the presence of a meiotically-derived embryo sac within the same ovule for endosperm formation.

In most plant species, the apomictic trait appears to be under the control of a single dominant locus. This locus may encode one or more developmental regulators, such as transcription factors, that in sexually reproducing plants function to initiate gene expression cascades leading to embryo sac and/or embryogenesis, but which are heterochronically or ectopically expressed in apomictic plants (Peacock, 1992; Koltunow, 1993; Koltunow et al, 1995).

Apomixis is a valuable trait for crop improvement since apomictic seeds give rise to clonal offspring and can therefore be used to genetically fix hybrid lines. The production of hybrid seed is a labour intensive and costly procedure as it involves maintaining populations of genetically pure parental lines, the use of separate pollen donor and male-sterile lines, and line isolation. Production of seed through apomixis avoids these problems in that once a hybrid has been produced, it can be maintained clonally, thereby eliminating the need to maintain and cross separate parental lines. The use of apomictic seed also provides a more cost effective method of multiplying vegetatively-propagated crops, as it eliminates the use of cuttings or tissue culture techniques to propagate lines, reduces the spread of diseases which are easily transmitted through vegetatively-propagated tissues, and in many species reduces the size of the propagule leading to lower shipping and planting costs.

Although apomixis occurs in a wide range of plant species, few crop species are apomictic. Attempts to introduce apomictic traits into crop species by introgression from wild relatives (Ozias-Akins, *et al.*, 1993; WO 97/10704; WO 97/11167) or through crosses between related, but developmentally divergent sexual species (WO 98/33374), have not yielded marketable products. Other approaches have focused on the identification of gene sequences that may be used to identify or manipulate apomictic

processes (WO 97/43427; WO 98/36090), however these approaches have not led to methods for the routine production of apomictic plants.

Mutagenesis approaches have also been attempted to convert sexually reproducing plants such as *Arabidopsis thaliana* (arabidopsis) into apomictic plants (Peacock *et al.*, 1995). For example, a number of recessive "fertilization-independent seed" (*fis*) mutants have been identified that initiate partial embryo and/or endosperm at a low frequency in the absence of fertilization (Chaudhury *et al.*, 1997). However, a number of additional parameters need to be modified in order to obtain true diploid apomictic seed using *fis* mutants.

Asexually-derived embryos can be induced to form in culture from many gametophytic and somatic plant tissues (Yeung, 1995). Somatic embryos can be obtained from culture of somatic tissues by treating them with plant growth regulators, such as auxins, or auxins in combination with cytokinins. Embryos can also be induced to form in culture from the gametophytic tissues of the ovule (gynogenesis) and the anther (androgenesis, pollen or microspore embryogenesis), either by the addition of plant growth regulators or by a simple stress treatment.

Several mutants have been identified that may be used to induce efficient production of embryos *in vitro*. These include recessive arabidopsis mutants with altered shoot meristems, for example *primordia timing (pt)*, *clavata (clv)1* and *clv3*, which were shown to enhance embryogenic callus formation when seedlings were germinated in the presence of auxin (Mordhorst *et al.*, 1998). The altered expression of two arabidopsis genes, *LEAFY COTYLEDON (LEC1)*; WO 98/37184, Lotan *et al.*, 1998) and *pickle*, have been shown to promote the production of somatic embryos in the absence of added growth regulators. The *LEC1* gene encodes a homologue of the HAP3 subunit of a CCAAT box-binding transcription factor (CBF). The *LEC1* gene controls many aspects of zygotic embryo development including desiccation tolerance and cotyledon identity. Ectopic over-expression of the *LEC1* gene in a *lec1* mutant background results in the production of 2 transgenic lines that occasionally form embryo-like structures on leaves.



- 5 -

These embryo-like structures express genes, such as those encoding seed storage proteins and oil body proteins, which are normally preferentially expressed in developing embryos. Plants containing a recessive mutant *PICKLE* gene produce a thickened, primary root meristem. Mutant *pickle* roots produce embryo-forming callus when the root tissue is separated from the rest of the plant and placed on minimal medium without growth regulators (Ogas *et al.*, 1997). Mutant *pickle* roots show morphological characteristics of developing seeds, such as oil bodies and, as with *LEC1* over-expressers, accumulate genes preferentially expressed in developing seeds.

Efficient production of apomictic seed is only likely to be realised through the identification and subsequent modification of developmental regulators, such as transcription factors, that are known to activate gene expression cascades leading to embryogenesis in both sexually-reproducing and apomictic plants. The present invention addresses this need by providing methods for the production of apomictic seeds comprising ectopic over-expression of an embryo-expressed AP2 domain containing transcription factor, BNM3.

---

## SUMMARY OF THE INVENTION

The present invention relates to asexual embryo formation and regeneration in plants. More specifically, it relates to processes for producing asexually-derived embryos, and for enhancing regeneration capacity in plants.

According to the present invention there is provided an isolated DNA molecule comprising a nucleotide sequence that hybridizes to SEQ ID NO:5 under stringent conditions, that comprises at least 23 contiguous nucleotides of SEQ ID NO:5, or that is at least 70% homologous with the nucleotide sequence defined by SEQ ID NO:5.

This invention further relates to an isolated DNA molecule that hybridizes to SEQ ID NO:5 under stringent conditions, comprising a nucleic acid sequence encoding a protein, wherein the protein when present at a sufficient level within a plant cell renders the cell embryogenic, increases the regenerative capacity of the plant cell, or both renders the cell embryogenic and increases the regenerative capacity of the plant cell. Included within the present invention is the above isolated DNA molecule comprising a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5 under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions. Also included within the present invention is a vector comprising the isolated DNA molecule as defined above, wherein the isolated DNA molecule is under control of a regulatory element that directs expression of said DNA in a plant cell. The regulatory element may be a constitutive, inducible, tissue specific or a developmental active, regulatory element.

This invention also embraces a transformed plant cell, a transformed plant, or seed obtained from a transformed plant, each comprising the vector as defined above

This invention relates to an isolated protein encoded by an isolated DNA molecule that hybridizes to the nucleotide sequence defined by SEQ ID NO:5 under stringent conditions, wherein the protein, when present at a sufficient level within a plant

- 7 -

cell renders the cell embryogenic, or increases the regenerative capacity of the plant cell. Also included is a protein encoded by an isolated DNA molecule that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5 under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions. This invention also embraces an isolated DNA molecule that encodes a protein as defined by SEQ ID NO:2 or SEQ ID NO:4. The invention also pertains to a protein comprising at least 70% homology with the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:4, or comprises from about 30 to about 541 amino acids of the sequence disclosed in SEQ ID NO:2, or comprises from about 30 to about 561 amino acids of the sequence disclosed in SEQ ID NO:4.

The present invention is also directed to a method of producing asexually derived embryos comprising:

- i) transforming a plant cell with a vector comprising an isolated DNA molecule that hybridizes to SEQ ID NO:5 under stringent conditions and which encodes a protein that when present at a sufficient level within the plant cell renders the plant cell embryogenic, or increases the regenerative capacity of the plant cell, or that comprises a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5 under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions;
- ii) ~~growing the plant cell to produce transformed tissue;~~
- iii) selecting the transformed tissue for occurrence of the isolated DNA molecule; and
- iv) assaying the transformed tissue for asexual embryo formation.

This invention also relates to the above method where the step of assaying (step iv)) involves assaying for somatic embryos, gametophytically-derived embryos, adventitious embryony, diplospory, or for haploid parthenogenesis of the embryo sac.

The present invention also embraces a method of producing an apomictic plant comprising:

- 8 -

- i) transforming a plant cell with a vector comprising an isolated DNA molecule that hybridizes to SEQ ID NO:5 under stringent conditions and which encodes a protein that when present at a sufficient level within said plant cell renders the plant cell embryogenic, or increases the regenerative capacity of the plant cell, or that comprises a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5 under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions;
- ii) selecting the transformed plant for occurrence of the isolated DNA molecule; and
- iii) assaying the transformed plant for asexual embryo formation.

This invention also relates to the above method where the step of assaying (step iii)) involves assaying for asexually-derived embryos, somatic embryos, gametophytically-derived embryos, adventitious embryony, diplospory, or for haploid parthenogenesis of the embryo sac.

The present invention is also directed to a method of producing asexually derived embryos comprising:

- i) transiently transforming a plant cell with a vector comprising an isolated DNA molecule that hybridizes to SEQ ID NO:5 under stringent conditions and which encodes a protein that when present at a sufficient level within the plant cell renders the plant cell embryogenic, or increases the regenerative capacity of the plant cell, or that comprises a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5 under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions;
- ii) growing the transiently transformed plant cell to produce transiently transformed tissue;
- iii) assaying the transiently transformed tissue for asexual embryo formation.

This invention is directed to the above method where the step of assaying (step iii)) involves assaying for asexually-derived embryos, somatic embryos, gametophytically-

- 9 -

derived embryos, adventitious embryony, diplospory, or for haploid parthenogenesis of the embryo sac.

The present invention also presents a method of modifying the regenerative capacity of a plant comprising

- i) transforming a plant cell with a vector comprising an isolated DNA molecule that hybridizes to SEQ ID NO:5 under stringent conditions and which encodes a protein that when present at a sufficient level within the plant cell renders the plant cell embryogenic, or increases the regenerative capacity of said plant cell, or that comprises a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5 under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions;
- ii) growing the transformed plant cell to produce transformed tissue; and
- iii) assaying the transformed tissue for enhanced regeneration as compared to wild type tissue.

This invention also embraces the above method wherein step iii) includes assaying in the absence of a growth regulator.

The present invention also relates to a method of modifying the regenerative capacity of a plant comprising:

- i) transiently transforming a plant cell with a vector comprising an isolated DNA molecule that hybridizes to SEQ ID NO:5 under stringent conditions and which encodes a protein that when present at a sufficient level within the plant cell renders the plant cell embryogenic, or increases the regenerative capacity of the plant cell, or that comprises a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5 under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions;

- 10 -

- ii) growing the transiently transformed plant cell to produce transiently transformed tissue;
- iii) assaying the transformed tissue for enhanced regeneration as compared to wild type tissue.

This invention also embraces the above method wherein step iii) includes assaying in the absence of a growth regulator.

The present invention also relates to a method of selecting a transformed plant comprising;

- i) transforming a normally non-regenerative plant with a vector comprising an isolated DNA molecule that hybridizes to SEQ ID NO:5 under stringent conditions and which encodes a protein that when present at a sufficient level within the plant cell renders the plant cell embryogenic, or increases the regenerative capacity of said plant cell, or that comprises a nucleotide sequence that hybridizes to nucleotides 1620- 4873 of SEQ ID NO:5 under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions; and
- ii) determining whether the transformed plant is able to regenerate under conditions in which the normally non-regenerative plant does not regenerate.

---

The present invention is also directed to an isolated DNA molecule comprising a DNA sequence that hybridizes to nucleotides 1-1619 of SEQ ID NO:5 under stringent conditions, or that comprises at least 22 contiguous nucleotides within nucleotides 1-1619 of SEQ ID NO:5. Also included within the scope of the present invention is a vector comprising the isolated DNA molecule as just defined, operably associated with a gene of interest, wherein the isolated DNA molecule directs the expression of the gene of interest within a plant cell. The gene of interest may be heterologous with respect to the isolated DNA molecule. The gene of interest may be selected from the group consisting of a pharmaceutically active protein, antibody, industrial enzyme, protein supplement, nutraceutical, storage protein, animal feed and animal feed supplement. This invention

- 11 -

also includes a transformed plant cell, a transformed plant, or seed obtained from the transformed plant, comprising the vector as just defined.

Furthermore, the present invention includes a method for directing the expression of a gene of interest within a developing embryo of a plant comprising transforming said plant with a vector containing an isolated DNA molecule that hybridizes to nucleotides 1-1619 of SEQ ID NO:5 under stringent conditions, or that comprises at least 22 contiguous nucleotides within nucleotides 1-1619 of SEQ ID NO:5.

This invention also pertains to a method of producing a protein of interest comprising

- i) transforming a plant with at least one vector, comprising an isolated DNA molecule that hybridizes to SEQ ID NO:5 under stringent conditions and which encodes a protein that when present at a sufficient level within the plant cell renders the plant cell embryogenic, or increases the regenerative capacity of said plant cell, or that comprises a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5 under stringent conditions, or to the nucleotide sequence as defined within SEQ ID NO:1 or 3 under stringent conditions to produce a transformed plant;
- ii) selecting the transformed plant for occurrence of the isolated DNA molecule; and
- ~~iii) growing the transformed plant in order to produce the protein of interest,~~  
wherein expression of the protein of interest is induced by the expression product of said isolated DNA.

This method may also comprise transforming the plant with a second vector comprising a nucleotide sequence encoding the protein of interest under the control of a regulatory element, wherein the regulatory element induced by the expression product of the isolated DNA. Furthermore, this method may also be used to produce a protein of interest wherein the protein of interest is a native protein.

- 12 -

## BRIEF DESCRIPTION OF THE DRAWINGS

These and other features of the invention will become more apparent from the following description in which reference is made to the appended drawings wherein:

**Figure 1** shows a schematic representation of the effect of culture temperature on the developmental fate of isolated microspores and pollen of *Brassica napus*. Late uninucleate microspores and early binucleate pollen cultured at 25 °C or lower continue to divide and form functional pollen grains (gametophytic), while the same microspores and pollen cultured at 32 °C undergo numerous sporophytic divisions, leading to the formation of haploid embryos (embryogenic). Late uninucleate microspores and early binucleate pollen cultured for one day at 25 °C, followed by culture at 32 °C may undergo gametophytic divisions, but form neither embryos nor mature pollen grains (non-embryogenic).

**Figure 2** shows the alignment of the DNA sequences depicted in SEQ ID NO:1 and SEQ ID NO:3. The ATG and TAG translation initiation and translation termination codons are shown in bold. Identical nucleotides are indicated by (\*) and gaps are indicated by (-).

**Figure 3** shows the alignment of the predicted protein sequences encoded by the DNA of SEQ ID NO:1 and SEQ ID NO:3. The amino acid sequence of the first AP2 domain repeat (repeat 1) and the second AP2 domain repeat (repeat 2), are shown in bold. Identical amino acids are indicated by an asterisk (\*) and mismatches by a dot (.) below the sequence alignment.

**Figure 4** shows the presence of two *BNM3* genes in the *Brassica napus* genome. A DNA gel blot containing restriction digests of *B. napus* c.v. Topas genomic DNA was hybridized to a *BNM3A* cDNA fragment under high stringency conditions. The *BNM3A* cDNA hybridizes to two DNA fragments under these conditions. These



- 13 -

fragments correspond to the *BNM3A* and *BNM3B* genes. The position of the molecular size markers (Lambda DNA *Hind* III restriction fragments) is indicated to the left the figure. The restriction enzymes used to digest the DNA are indicated above the blot.

Figure 5 shows the alignment of the predicted protein sequence encoded by the DNA of SEQ ID NO.1 (*BNM3A*) with the predicted protein sequences of other AP2 domain proteins. The amino acid sequence of *BNM3A*, beginning at position 208, and spanning the first AP2 domain repeat (AP2 domain repeat 1), the second AP2 domain repeat (AP2 domain repeat 2), and the linker region lying between the two repeats (linker), was aligned with the amino acid sequence of other proteins containing two AP2 domains. The amino acid similarity in this region ranges from 53% for *APETALA2* to 80% for *ZMMHCF1*. Identical amino acids are indicated by (\*) and gaps are indicated by (-). Protein names are indicated on the left and are abbreviated as follows: ANT, *AINTEGUMENTA* (accession number U41339); ZM, *ZMMHCF1* (accession number Z47554); GL15, *GLOSSY15* (accession number U41466); AP2, *APETALA2* (accession number U12546).

Figure 6 shows the results of gel blot analysis with a *BNM3A* cDNA fragment performed on RNA extracted from the indicated tissues. RNA gel blots contain either 5 µg (a) or 20 µg (b, c) of total RNA. Figure 6A shows the pattern of *BNM3* expression in microspore-embryo cultures. RNA was isolated from late uninucleate microspores and early binucleate pollen at the time of collection (pollen 0d), after four days in culture at 32 °C (+ embryo), after four days in culture at 25 °C (pollen 4d), after one day of culture at 25 °C, followed by three days of culture at 32 °C (- embryo) and microspore-derived embryos at the globular, heart, torpedo, 21 day old cotyledon (21 d cot), 28 day old cotyledon (28 d cot) and 42 day old cotyledon (42 d cot) stage of development. *BNM3* expression is detected in embryogenic microspores and developing microspore-derived embryos, but is absent from developing microspores and pollen collected prior to tissue culture and in non-embryogenic samples. The exposure time was seven days. Figure 6B

- 14 -

shows that *BNM3* gene expression is detected in developing seeds. Seeds were collected at various days after pollination (DAP). These points in development correspond approximately to the globular (7 d), heart (14 d), torpedo (18 d), early cotyledon (21 d), mid cotyledon (28 d, 35 d) and late cotyledon (42 d) stages of development. The exposure time was 14 days. Figure 6C shows that *BNM3* gene expression is not detected in non-seed tissues. Roots and leaves were collected from 14 day old greenhouse grown plants. Entire flowers as well as excised anthers and pistils were collected from opened flower buds just prior to anthesis. Small and large buds refer to closed flower buds of less than 5 mm or greater than 5 mm in length, respectively. Siliques were collected 16 days after pollination. The exposure time was 14 days.

Figure 7 shows the phenotype of *Brassica napus* and arabidopsis plants transformed with constructs containing the *BNM3* gene under control of a modified *POLYUBIQUITIN* promoter (B) and double enhanced 35S promoter containing an AMV translational enhancer (A, C-E). Figure 7A shows embryo structures on the leaf margin of a *Brassica* T1 seedling. Figure 7B shows embryo structures on the petiole of an arabidopsis T2 seedling. Figure 7C shows embryo structures on the cotyledon of an arabidopsis T1 seedling. Figure 7D shows a scanning electron micrograph of the abaxial side of an arabidopsis T1 cotyledon. Note the bipolar nature of the embryos, as well as the emergence of a secondary embryo from the surface of a primary embryo (asterisk). Figure 7E shows a semi-thin section through one of the cotyledons of the T1 seedling shown in (Figure 7C). Note the presence of all the major organs and tissue elements of embryo, as well as the development of new embryos on the flanks of the shoot apical meristems and the cotyledons.

Figure 8 shows the increased regenerative capacity of arabidopsis plants transformed with a construct containing the *BNM3B* gene under control of a modified *POLYUBIQUITIN* promoter. Figure 8A shows wild-type and transgenic leaf and hypocotyl explants on medium containing growth regulators. Figure 8B shows

- 15 -

wild-type and transgenic roots on medium containing growth regulators. Figure 8C shows wild-type and transgenic leaf and hypocotyl explants on medium without growth regulators. Figure 8D shows wild-type and transgenic root explants on medium without growth regulators

- 16 -

**DESCRIPTION OF PREFERRED EMBODIMENT**

The present invention relates to asexual embryo formation and regeneration in plants. More specifically, it relates to processes for producing asexually-derived embryos, and for enhancing regeneration capacity in plants. The present invention also relates to heterologous protein production systems in plants, and the uses thereof.

Genes preferentially expressed during the induction of *Brassica napus* c.v. Topas microspore embryogenesis were isolated via subtractive screening. Seven independent cDNA clones, comprising six unique DNA sequences were found to be differentially expressed between cDNA libraries prepared from embryogenic and non-embryogenic microspore cultures. Several of these *BNM* (for *Brassica napus* microspore embryo) clones, *BNM3A* (SEQ ID NO:1) and *BNM3B* (SEQ ID NO:3), were characterized as described herein. *BNM3A* and *BNM3B* encode the amino acid sequences disclosed in SEQ ID NO:2, and SEQ ID NO:4, respectively. The genomic sequence of *BNM3A* (SEQ ID NO:5), including the regulatory region (nucleotides 1-1619 of SEQ ID NO:5), was also obtained.

"Regeneration", as used herein, refers to a morphogenetic response that results in the production of new tissues, organs, embryos, whole plants or fragments of whole plants that are derived from a single cell, or a group of cells. Regeneration may proceed ~~indirectly via a callus phase or directly, without an intervening callus phase~~. "Regenerative capacity" refers to the ability of a plant cell to undergo regeneration.

By "embryogenic cell", it is meant a cell that has completed the transition from either a somatic or a gametophytic cell to a state where no further applied stimuli are necessary to produce a somatic or gametophytic embryo, respectively.

By "regulatory element" it is meant those that include developmentally regulated, tissue specific, inducible and constitutive regulatory elements. A regulatory element that is developmentally regulated, or controls the differential expression of a gene under its

- 17 -

control, is activated within certain organs or tissues of an organ at specific times during the development of that organ or tissue. However, some regulatory elements that are developmentally regulated may preferentially be active within certain organs or tissues at specific developmental stages, they may also be active in a developmentally regulated manner, or at a basal level in other organs or tissues within the plant as well, such regulatory elements are considered "tissue specific". Regulatory elements may be found either upstream, within, downstream, or a combination thereof, of the coding region of a gene.

An inducible regulatory element is one that is capable of directly or indirectly activating transcription of one or more DNA sequences or genes in response to an inducer. In the absence of an inducer the DNA sequences or genes will not be transcribed. Typically the protein factor, that binds specifically to an inducible regulatory element to activate transcription, is present in an inactive form which is then directly or indirectly converted to the active form by the inducer. The inducer can be a chemical agent such as a protein, metabolite, growth regulator, herbicide or phenolic compound or a physiological stress imposed directly by heat, cold, salt, or toxic elements or indirectly through the action of a pathogen or disease agent such as a virus. A plant cell containing an inducible regulatory element may be exposed to an inducer by externally applying the inducer to the cell or plant such as by spraying, watering, heating or similar methods.

A constitutive regulatory element directs the expression of a gene throughout the ~~various parts of a plant and continuously throughout plant development. Examples of~~ known constitutive regulatory elements include promoters associated with the CaMV 35S transcript. (Odell et al., 1985, *Nature*, **313**: 810-812), the rice actin 1 (Zhang et al, 1991, *Plant Cell*, **3**: 1155-1165) and triosephosphate isomerase 1 (Xu et al, 1994, *Plant Physiol.* **106**: 459-467) genes, the maize ubiquitin 1 gene (Corno et al, 1993, *Plant Mol. Biol.* **29**: 637-646), the *Arabidopsis* ubiquitin 1 and 6 genes (Holtorf et al, 1995, *Plant Mol. Biol.* **29**: 637-646), and the tobacco translational initiation factor 4A gene (Mandel et al, 1995 *Plant Mol. Biol.* **29**: 995-1004).

By "gene of interest" it is meant any gene that is to be expressed in a transformed plant. Such a gene of interest may include, but is not limited to, a gene that encodes a pharmaceutically active protein, for example growth factors, growth regulators, antibodies, antigens, their derivatives useful for immunization or vaccination and the like. Such proteins include, but are not limited to, interleukins, insulin, G-CSF, GM-CSF, hPG-CSF, M-CSF or combinations thereof, interferons, for example, interferon- $\alpha$ , interferon- $\beta$ , interferon- $\gamma$ , blood clotting factors, for example, Factor VIII, Factor IX, or tPA or combinations thereof. A gene of interest may also encode an industrial enzyme, protein supplement, nutraceutical, or a value-added product for feed, food, or both feed and food use. Examples of such proteins include, but are not limited to proteases, oxidases, phytases chitinases, invertases, lipases, cellulases, xylanases, enzymes involved in oil biosynthesis etc. Other protein supplements, nutraceuticals, or a value-added products include native or modified seed storage proteins and the like.

The present invention is further directed to a chimeric gene construct containing a DNA of interest operatively linked to a regulatory element of the present invention. Any exogenous gene, or gene of interest, can be used and manipulated according to the present invention to result in the expression of the exogenous gene.

The activation of the expression of a gene of interest may also be under the control of a regulatory element that itself is activated by a BNM3 protein. For example, ~~which is not to be considered limiting, a gene of interest may be fused to the napin promoter, and the napin promoter may be induced by BNM3.~~ Furthermore, a gene of interest may be expressed within somatic tissues under the control of one or more regulatory elements induced by BNM3, so that, as will be described in more detail below, the somatic tissue develops into a seed-like structure comprising embryogenic cells, and these seed-like structures produce the products of the gene of interest.

The chimeric gene construct of the present invention can further comprise a 3' untranslated region. A 3' untranslated region refers to that portion of a gene comprising a DNA segment that contains a polyadenylation signal and any other regulatory signals

- 19 -

capable of effecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by effecting the addition of polyadenylic acid tracks to the 3' end of the mRNA precursor. Polyadenylation signals are commonly recognized by the presence of homology to the canonical form 5' AATAAA-3' although variations are not uncommon. Examples of suitable 3' regions are the 3' transcribed non-translated regions containing a polyadenylation signal of *Agrobacterium* tumor inducing (Ti) plasmid genes, such as the nopaline synthase (*Nos* gene) and plant genes such as the soybean storage protein genes and the small subunit of the ribulose-1, 5-bisphosphate carboxylase (ssRUBISCO) gene. The 3' untranslated region from the structural gene of the present construct can therefore be used to construct chimeric genes for expression in plants.

The chimeric gene construct of the present invention can also include further enhancers, either translation or transcription enhancers, as may be required. These enhancer regions are well known to persons skilled in the art, and can include the ATG initiation codon and adjacent sequences. The initiation codon must be in phase with the reading frame of the coding sequence to ensure translation of the entire sequence. The translation control signals and initiation codons can be from a variety of origins, both natural and synthetic. Translational initiation regions may be provided from the source of the transcriptional initiation region, or from the structural gene. The sequence can also be derived from the regulatory element selected to express the gene, and can be specifically modified so as to increase translation of the mRNA.

---

To aid in identification of transformed plant cells, the constructs of this invention may be further manipulated to include plant selectable markers. Useful selectable markers include enzymes which provide for resistance to an antibiotic such as gentamycin, hygromycin, kanamycin, and the like. Similarly, enzymes providing for production of a compound identifiable by colour change such as *GUS* ( $\beta$ -glucuronidase), fluorescence, or luminescence, such as luciferase are useful.

Also considered part of this invention are transgenic plants containing a gene or chimeric gene construct of the present invention comprising a *BNM3* gene, a regulatory

- 20 -

element obtained from *BNM3*, or the coding region from *BNM3* in operative association with a constitutive, developmental or inducible regulatory element, or a combination thereof. Methods of regenerating whole plants from plant cells are known in the art. In general, transformed plant cells are cultured in an appropriate medium, which may contain selective agents such as antibiotics, where selectable markers are used to facilitate identification of transformed plant cells. Once callus forms, shoot formation can be encouraged by employing the appropriate plant hormones in accordance with known methods and the shoots transferred to rooting medium for regeneration of plants. The plants may then be used to establish repetitive generations, either from seeds or using vegetative propagation techniques. The constructs of the present invention can be introduced into plant cells using Ti plasmids, Ri plasmids, plant virus vectors, direct DNA transformation, micro-injection, electroporation, biolistics etc. For reviews of such techniques see for example Weissbach and Weissbach, *Methods for Plant Molecular Biology*, Academy Press, New York VIII, pp. 421-463 (1988); Geierman and Corey, *Plant Molecular Biology*, 2d Ed. (1988); and Miki and Iyer, *Fundamentals of Gene Transfer in Plants*. In *Plant Metabolism*, 2d Ed. DT. Dennis, DH Turpin, DD Lefebvre, DB Layzell (eds), Addison Wesley, Langmans Ltd. London, pp. 561-579 (1997). The present invention further includes a suitable vector comprising the gene or the chimeric gene construct.

A class of genes have been isolated from *Brassica napus* microspore embryo cultures. These genes have been found to be important regulators of embryogenesis by ~~their ability to induce the formation of asexually-derived embryos when ectopically~~ expressed in the vegetative tissues of plants. These genes are hereinafter indicated as *BNM3* genes (*Brassica napus* microspore embryo). SEQ ID NO. 1 depicts the cDNA of *BNM3A*, SEQ ID NO. 3 depicts the cDNA of *BNM3B*, and the genomic sequence for *BNM3A* is given in SEQ ID NO:5. The promoter of *BNM3A* lies within nucleotides 1-1619 of SEQ ID NO:5. The predicted protein sequences encoded by the DNAs of SEQ ID NO. 1 and 3 are outlined in SEQ ID NOs. 2 and 4, respectively.

By "*BNM3*" or "*BNM3* gene", it is meant the sequence of oligonucleotides as disclosed in SEQ ID NOs:1, 3 or 5, or fragments, derivatives, or mutations thereof, or oligonucleotide sequences that exhibit at least 70% homology or similarity, with a



- 21 -

fragment or derivative of the sequences disclosed in SEQ ID NOs 1, 3, or the coding region (nucleotides 1620-4873) of SEQ ID NO:5, as determined using oligonucleotide alignment algorithms (for example, but not limited to a BLAST or FASTA). Furthermore, oligonucleotides that associate with these sequences under conditions of high stringency, for example, but not to be limited to, hybridization to gel blots at about 65°C followed by wash conditions at 0.1X SSC, 65°C, are also considered *BNM3* genes. "*BNM3* gene" also includes DNA molecules that comprises at least 23 contiguous nucleotides of SEQ ID NOs:1, 3 or 5, or at least 22 contiguous nucleotides within nucleotides 1-1619 of SEQ ID NO:5. A fragment of *BNM3*, that comprises at least 22 contiguous nucleotides may be used as a probe for the identification of nucleotides related to *BNM3* regulatory, or coding, regions within an organism. Furthermore, molecules comprising at least 23 contiguous nucleotides of SEQ ID NOs:1, 3 or 5 and that encode a protein, or an active fragment thereof, that when present at a sufficient level within a plant cell renders the cell embryogenic, increases the regenerative capacity of the plant cell, or renders the cell embryogenic and increases the regenerative capacity of the plant cell are also considered to be *BNM3* genes. Preferably, a *BNM3* gene comprises from about 50 to about 1981 nucleotides of SEQ ID NOs: 1 or 3, or about 50 to about 4858 nucleotides from the coding region (1620-4858) of SEQ ID NO:5.

By "*BNM3* regulatory region" it is meant the sequence of oligonucleotides from 1-1619 in SEQ ID NO:5, or a fragment, derivative, or mutation thereof. Furthermore, a ~~*BNM3* regulatory region also comprises a nucleotide sequence that associates with a~~ sequence from 1-1619 of SEQ ID NO:5 under conditions of high stringency, for example, but not to be limited to, hybridization to gel blots at about 65°C followed by wash conditions at 0.1X SSC, 65°C, or that exhibits at least 70% homology or similarity, with a fragment or derivative of the sequences disclosed in nucleotides 1-1619 of SEQ ID NO:5, as determined using oligonucleotide alignment (for example, but not limited to a BLAST or FASTA search).

By "*BNM3* protein" it is meant a protein, or a biologically active fragment thereof, that renders a plant cell embryogenic, increases the regenerative capacity of the

- 22 -

plant cell, or renders the cell embryogenic, increases the regenerative capacity of the plant cell, and that is encoded by a *BNM3* gene, as defined above. Preferably, a *BNM3* protein comprises from about 30 to about 541 amino acids of the sequence disclosed in SEQ ID NO:2, or from about 30 to about 561 amino acids of the sequence disclosed in SEQ ID NO: 4. However, *BNM3* protein may also be defined as a protein having at least 70% homology with either SEQ ID NO:2 or 4.

Search of the sequence databases indicated that the *BNM3* translation products contain two copies of an AP2 domain (Figure 3; see also SEQ ID NO: 2 for *BNM3A*, and SEQ ID NO: 4 for *BNM3B*). The AP2 domain was first identified in *APETALA2*, an arabidopsis protein that regulates meristem identity, floral organ specification, seedcoat development and floral homeotic gene expression (Jofuku *et al.*, 1994), but has since been identified in a wide range of proteins with diverse functions.

The AP2 domain is usually between 58 to 68 amino acids in length and contains a conserved central core of 18 amino acids, characterized by its ability to form an amphipathic  $\alpha$  helix, a structure thought to mediate protein-protein interactions. The ability of a number of AP2 domain containing proteins to bind DNA, coupled with the presence of putative nuclear localization signals and acidic regions that may function as transcriptional activators suggests these proteins function as transcription factors.

---

Two phylogenetically distinct classes of AP2 domain proteins have been identified; proteins with a single AP2 domain (EREBP-like) and proteins with two AP2 domains (AP2-like; (Zhou, 1997)). The proteins encoded by the genes of this invention represent unique members of the latter class of proteins.

---

Accordingly, an aspect of the present invention provides for an isolated DNA molecule that comprises a sequence encoding a protein that contains two AP2 domains. The protein, when present at a sufficient level in a plant cell, renders the cell embryogenic, increases the regenerative capacity of the cell, or both renders the cell embryogenic and increases the regenerative capacity of the cell.

Analysis of *BNM3* expression during microspore-derived embryo development, seed development, or non-seed tissue development, using Northern (Figure 6) indicated that the *BNM3* genes are preferentially expressed in embryogenic microspore cultures, microspore-derived embryos and seeds. *BNM3* transcripts were not detected in any of the non-seed tissues tested.

*BNM3* mRNA is detected in microspore cultures induced to undergo embryogenesis, as well as in the subsequent globular, heart, torpedo and cotyledon stages of microspore-derived embryo development (e.g. Figure 6A). RNAs are also detected within developing seeds, 14 days after pollination (14 DAP), corresponding to the heart stage of embryo development. *BNM3* expression increases during the early (21 DAP) and mid-cotyledon (28 DAP) stages of embryo development and remains constant thereafter (Figure 6B).

Constitutive expression of *BNM3* resulted in the formation of somatic embryos on vegetative structures such as cotyledons, petioles, leaf blades and the shoot apical meristem of plants (Figure 7). In these experiments *BNM3* cDNAs were placed under the control of two separate constitutive promoter constructs, a modified sunflower *POLYUBIQUITIN* promoter construct, and a double enhanced 35S promoter construct containing an AMV translational enhancer, however, it is to be understood that any suitable constitutive promoter may be used for this purpose. Such *BNM3*-derived ectopic embryos contain all of the organ systems and tissue layers found in the developing zygotic embryo in that these embryos are bipolar (Figure 7E), consist of an axis, a hypocotyl and radicle region, shoot and root meristems, and cotyledons. In addition, each organ system contained the characteristic radial arrangement of three specialized tissue layers (epidermis, ground parenchyma and provascular tissue) found in zygotic embryos. Continued expression of the *BNM3* gene within the developing ectopic embryo leads to a reiteration of the embryo-forming process, with the result that new embryos are continuously formed on the surface of pre-existing embryos (Figure 7E).

- 24 -

Constitutive expression of *BNM3* results in the increased ability of a plant to regenerate shoots *in vitro* in the presence of added growth regulators. Root explants from transgenic plants ectopically expressing *BNM3* show at least a 5-fold increase in shoot regeneration in the presence of hormones as compared to root explants obtained from wild-type plants (Figure 8A,B). Shoots also developed faster in the transgenic explants, compared to the wild-type. Wild-type leaf and hypocotyl explants initially responded by producing callus on the cut end of the petiole (Figure 8B) followed by callus formation along the length of the petiole. In contrast, explants from transgenic lines immediately produced new shoots (Figure 8B) or roots from the cut end of the petiole. Explants that initially produced roots eventually also produced shoots.

Transgenic explants, constitutively expressing *BNM3* were also able to regenerate in the absence of added growth regulators. These explants, when placed on media lacking growth regulators regenerated shoots either from the cut end of the leaf and hypocotyl explants or from the nodule-like structures of root explants (Figures 8C,D). In all cases regenerated shoots developed, rooted, flowered and set seed. Conversely, wild-type leaf and hypocotyl explants placed on medium lacking growth regulators occasionally produce callus or roots at the cut end of the leaf petiole, however no shoots form from these structures (Figure 8C,D).

It is also considered within the scope of the present invention, that expression of ~~*BNM3* may be used to initiate a developmental cascade within a transformed plant or~~ plant cell. This cascade may arise as a result of the stable integration of a DNA-based vector expressing *BNM3* within a transformed plant, however, such a cascade may also arise as a result of transient expression of *BNM3*, and does not require the stable integration of the *BNM3*-based vector within a plant cell. These transient approaches may be useful for inducing somatic embryogenesis, gametophytically-derived embryogenesis, or increasing the regenerative capacity of a plant or plant cell.

Plants in which a *BNM3* gene is ectopically expressed exhibit advantageous qualities including:

- 25 -

- formation of asexually derived embryos;
- increased regenerative capacity of tissue explants;
- the ability of tissue explants to regenerate in the absence of added plant growth regulators; and
- the expression of seed components in non-seed organs in which *BNM3* is ectopically expressed.

Furthermore, plants that ectopically express at least one *BNM3* gene can be used for the production of recombinant proteins using seed specific regulatory elements.

For the applications of *BNM3* as described below, it will be advantageous to obtain a high level of the *BNM3* transcript and/or *BNM3* protein in order to obtain plants in which the phenotype is highly penetrant. This may be obtained by using genetic elements such as introns, transcriptional enhancers or translational enhancers which are known to enhance gene or protein expression levels.

The *BNM3* sequences of the present invention may be used for several applications including, but not limited to, the control of embryo processes, the control of regeneration processes, the use of regulatory sequences for targeted gene expression, the use of *BNM3* sequences as selectable markers of transformed plants, or for embryogenic cells. These applications are disclosed in more detail below.

---

#### Use of *BNM3* Sequences to Control Embryogenic Processes

---

As described herein, *BNM3* genes play an important role in initiation and maintenance of embryo development. *BNM3* genes have been found in a wide range of members of the plant kingdom. Regulatory regions obtained from these genes may be used to control the transcription of *BNM3* or a derivative or fragment thereof, or any gene of interest, using methods known to one of skill in the art.

Ectopic expression of a *BNM3* gene is sufficient to induce recurrent formation of asexually derived embryos on the vegetative tissues of plants (see example 4). Depending

- 26 -

upon the promoter used, ectopic over-expression of *BNM3* genes may be used to produce somatic or gametophytic embryos. Somatic or gametophytic embryos may be obtained by expressing a *BNM3* gene under the control of a constitutive regulatory element, as is shown in Example 5, or may also be obtained by expressing a *BNM3* gene under the control of tissue specific or developmentally regulated elements, inducible elements derived from either plant or non-plant genes or through transient expression. In this respect, chemical induction systems (e.g. see Gatz and Lenk, 1998, which is incorporated by reference) or transient expression using methods which do not result in stable integration of the *BNM3* gene, or which make direct use of the *BNM3* protein e.g. microprojectile bombardment of DNA or protein may also be employed.

Temporal and/or spatial restriction of *BNM3* expression using inducible, tissue specific or developmentally regulated elements, is preferred when recurrent embryogenesis is not a desirable trait. The regulatory elements used to restrict *BNM3* to a specific developmental stage or cell type will depend on the application. For example, regulatory elements that may be used to express *BNM3* for the production of microspore-derived embryos include, but are not limited to, those of the class I low molecular weight heat shock inducible gene, *GMHSP17.3B* (Zarsky *et al.*, 1995, which is incorporated by reference), or microspore/pollen expressed genes such as *NTM19* (Custers *et al.*, 1997, EP 790,311, which are incorporated by reference), *BCP1* (Xu *et al.*, 1995, which is incorporated by reference), *LATS2* (Twell *et al.*, 1989, which is incorporated by reference), *BNM1* (Treacy *et al.* 1997, which is incorporated by reference), and *APG* (Roberts *et al.*, 1993, which is incorporated by reference).

Examples of regulatory elements that may be used to express *BNM3* for the production of somatic embryos include, but are not limited to, those of genes activated by plant growth regulators which are routinely used to induce somatic embryogenesis in tissue culture. Specific examples, which are to be considered non-limiting, include the cytokinin inducible *IB6* and *CKII* genes (Brandstatter and Kieber, 1998; Kakimoto, 1996, which are incorporated by reference) and the auxin inducible element, DR5 (Ulmasov *et*

- 27 -

*al.*, 1997, which is incorporated by reference). However, it is to be understood that other regulatory elements may be included for the expression of *BNM3* in plants.

Furthermore, examples of gene regulatory elements suitable for directing expression of *BNM3* to obtain adventitious embryony include, but are not limited to, those obtained from the ovule and embryo expressed *SERK* gene (Schmidt *et al.*, 1997 which is incorporated by reference), the ovule expressed *AGL11* gene (Roundsley *et al.*, 1995, which is incorporated by reference), the nucellus expressed *NUC1* gene (Doan *et al.*, 1996; WO 98/08961, which are incorporated by reference), or the inner integument-expressed genes, *FBP7* (Angenent *et al.*, 1995, which is incorporated by reference) and *SC4* (US application 09/059,909, filed April 13, 1998, which is incorporated by reference) genes.

According to one aspect of the present invention there is provided a method for the efficient production of microspore-derived embryos in plants. This method involves:

- i) transforming a plant of interest, for example, *Brassica napus* (using transformation techniques known to one of skill, for example, DeBlock *et al.*, 1989, Clough and Bent 1998, Vergunst *et al.* 1998, Klein *et al.* 1987, which are incorporated herein by reference) with a vector construct, or isolated DNA, consisting of a *BNM3* gene under control of a suitable regulatory element, which ~~may be constitutive, tissue specific, developmentally regulated, or inducible and~~ optionally, a marker gene for selection of transformants;
- ii) selecting transformed plants;
- iii) producing lines that ectopically overexpress the *BNM3* gene, or *BNM3* protein;
- iv) isolating microspores and pollen from the transgenic lines and culturing microspores and pollen to induce embryogenesis.

Embryogenesis can be induced by any suitable protocol, for example, which is not to be considered limiting, culturing microspore and pollen for about four days at from about 28° to about 35 °C, preferably at about 32°C, then transferring embryogenic cells or embryos to about 25 °C.

- 28 -

Using the above method, *Brassica napus* cultivars ectopically overexpressing *BNM3* show an increase in the percentage of embryogenic cells or embryos over that observed when microspores or pollen are prepared from wild-type plants that do not ectopically express *BNM3*.

Examples of regulatory elements that may be used to express *BNM3* for the production of microspore-derived embryos include, but are not limited to, those of the class I low molecular weight heat shock inducible gene, *GMHSP17.3B* (Zarsky *et al.*, 1995, which is incorporated by reference), or microspore/pollen expressed genes such as *NTM19* (Oldenhof *et al.*, 1996, EP 790,311, which are incorporated by reference), *BCP1* (Xu *et al.*, 1995, which is incorporated by reference), *LAT52* (Twell *et al.*, 1989, which is incorporated by reference), *BNM1* (Treacy *et al.* 1997, which is incorporated by reference), and *APG* (Roberts *et al.*, 1993, which is incorporated by reference). Also useful are inducible regulatory elements, for example but not limited to, tetracycline-inducible promoter (Gatz 1997, which is incorporated by reference), steroid inducible promoter (Aoyama and Chua 1997, which is incorporated by reference) and ethanol-inducible promoter (Slater *et al.* 1998, Caddick *et al.* 1998, which are incorporated by reference).

In a similar fashion, microspore-derived embryos may also be produced in plants by introducing into a plant of interest a *BNM3* protein. (e.g. via biolistics : Klein *et al.* 1987) and selecting for plants that exhibit increased microspore embryogenesis.

This invention also provides a method for the efficient production of somatic embryos *in vitro*. This method involves:

- i) transforming a plant, for example, *Arabidopsis* using transformation techniques known to one of skill (for example, but not limited to, DeBlock *et al.*, 1989, Clough and Bent 1998, Vergunst *et al.* 1998, which are incorporated by reference), or a plant cell may also be transiently transformed using methods



- 29 -

known to one of skill (for example, biolistics; Klein et al 1987) with a vector construct containing a *BNM3* gene under control of suitable regulatory element, which may be constitutive, inducible or developmentally regulated, and, optionally, a marker gene for selection of transformants is transformed to several arabidopsis.

- ii) selecting transformed plants, and
- iii) culturing the desired explant from the selected transformed plants, for example, but not limited to, root, leaf or seedlings *in vitro*, in media with or without appropriate growth regulators, for example, but not limited to 2,4-D (e.g. Mordhorst *et al.*, 1998) to produce direct embryogenesis or embryogenic callus; and
- iv) transferring embryos, non-embryogenic callus, or both embryos and non-embryogenic callus to appropriate media for the production of embryos, plantlets, or both embryos or plantlets.

For example, when the results of the above method are compared with the production of somatic embryos *in vitro* using a number of *Arabidopsis* ecotypes, directed embryogenesis or embryogenic callus is initiated at a higher frequency from transgenic lines ectopically over-expressing *BNM3* than in wild-type controls.

Examples of regulatory elements that may be used to express *BNM3* for the production of somatic embryos include, but are not limited to, those of genes activated by plant growth regulators which are routinely used to induce somatic embryogenesis in tissue culture. Specific examples, which are to be considered non-limiting, cytokinin inducible *IB6* and *CK11* genes (Brandstatter and Kieber, 1998; Kakimoto, 1996, which are incorporated by reference) and the auxin inducible element, DR5 (Ulmasov *et al.*, 1997, which is incorporated by reference). Also useful are inducible regulatory elements, for example but not limited to, a tetracycline-inducible promoter (Gatz 1997, which is incorporated by reference), a steroid inducible promoter (Aoyama and Chua 1997, which is incorporated by reference), and an ethanol-inducible promoter (Slater et al 1998, Caddick et al. 1998, which are incorporated by reference).

- 30 -

Ectopic initiation of embryo development is one of the key steps in apomixis. As shown in Example 4, ectopic expression of a *BNM3* gene is sufficient to initiate embryo formation in otherwise non-embryo-forming tissue. A *BNM3* gene may therefore be used to initiate adventitious embryony or parthenogenesis of a reduced or unreduced embryo sac cell by expression of the gene in the sporophytic or gametophytic tissues of the developing ovule.

Adventitious embryony is achieved by expressing *BNM3* in sporophytic ovule tissues such as the nucellus, the inner integuments or other tissues lying adjacent to or in proximity to the developing embryo sac. This method involves:

- i) transforming a desired plant (see above methods) with a vector construct consisting of a *BNM3* gene under control of suitable regulatory element, which may be constitutive, inducible or developmentally regulated, and, optionally, a marker gene for selection of transformants, using methods known within the art;
- ii) selecting transformed plants;
- iii) emasculating the transformed plant;
- iv) pollinating the transformed plants with pollen carrying one or more dominant selectable markers, for example GUS or kanamycin resistance; and
- v) assaying for production of clonal offspring.

When the results of the above method are compared with the pollination of a wild-type ~~arabidopsis plant with pollen carrying the dominant selectable marker, all F1 embryos~~ resulting from this cross inherit the dominant marker while embryos derived from plants ectopically over expressing the *BNM3* gene or protein are clonally derived via sexual embryo formation and do not inherit the dominant selectable marker.

Specific examples of gene regulatory elements suitable for directing expression of *BNM3* to obtain adventitious embryony, diplospory or haploid parthenogenesis of embryo sac components include the ovule expressed SERK gene (Schmidt et al. 1997, which is incorporated by reference), the meiosis expressed *AtDMC1* gene, (Klimyuk and Jones, 1997; WO 98/28431, which are incorporated by reference), the ovule expressed

*AGL11* gene (Roundsley *et al.*, 1995, which is incorporated by reference), the nucellus expressed *NUC1* gene (Doan *et al.*, 1996; WO 98/08961, which are incorporated by reference), and the inner integument-expressed genes, *FBP7* (Angenent *et al.*, 1995, which is incorporated by reference) and *SC4* (US application 09/059,909, filed April 13, 1998, which is incorporated by reference) genes. Furthermore, inducible systems, for example but not limited to, tetracycline-inducible promoter (Gatz 1997, which is incorporated by reference), steroid inducible promoter (Aoyama and Chua 1997, which is incorporated by reference), ethanol-inducible promoter (Slater *et al.* 1998, Caddick *et al.* 1998, which are incorporated by reference) may also be used. Parthenogenesis from cells of the embryo sac requires a regulatory element that is active in one or more cells of the female gametophyte or their precursors. Fertilization of the meiotically-derived polar nuclei is desirable when the development of seed is dependent on the presence of endosperm.

#### Use of *BNM3* Sequences to Control Regeneration Processes

Plants ectopically over-expressing the *BNM3* genes exhibit increased regenerative capacity and the ability to regenerate whole plants in the absence of added growth regulators (see example 5). *BNM3* gene expression may therefore be used to enhance or induce the regeneration capacity of plant tissues *in vivo* or *in vitro*. The regulatory elements used to express *BNM3* will depend, in part, on the target tissue used for regeneration. Regeneration of plant tissues may be obtained by

expressing a *BNM3* gene under the control of a constitutive regulatory element, for example, but not limited to, 35S, or by expressing a *BNM3* gene under the control of tissue specific or developmentally regulated elements, inducible elements derived from either plant or non-plant genes (e.g. Gatz and Lenk, 1998, which is incorporated by reference), or through transient expression methods which do not result in stable integration of the *BNM3* gene or which make direct use of the *BNM3* protein (e.g. microprojectile bombardment of DNA or protein). Chemical induction systems (see Gatz and Lenk, 1998) or regulatory elements of genes that respond to plant growth regulators used to induce regeneration, such as, for example, cytokinin (Brandstatter

- 32 -

and Kieber, 1998; Kakimoto, 1996) or auxin (Ulmasov et al., 1997), or genes expressed at the wound site of tissue explants (Xu *et al.*, 1993) may be used.

A further application is the use of a *BNM3* gene as a selectable marker for the recovery of transgenic plants. As an example of this application which is not to be considered limiting in any manner, roots of a seedling, for example, an *Arabidopsis* ecotype C24 seedling, are cocultivated with a single *Agrobacterium tumefaciens* strain (per Vergunst *et al.*, 1998; except that all steps are carried out in the absence of added growth regulators) containing two binary constructs:

- a first binary vector carries a reporter gene fusion, for example, but not limited to, *35S::GUS*;
- a second binary vector contains a *BNM3* gene under control of suitable regulatory element.

*BNM3* gene expression is activated upon integration of the above construct into the arabidopsis genome and transgenic plants are selected on the basis of their ability to regenerate under conditions in which wild-type explants are unable to regenerate, for example, but not limited to, the absence of growth regulators. In many instances the T-DNA carrying the *BNM3* gene and the T-DNA carrying the gene of interest will integrate at unlinked loci. The T-DNA containing the introduced *BNM3* sequence, and its associated increased regenerative capacity phenotype, may therefore be removed in the progeny plants by simple segregation (Daley et al. 1998). However, as will be apparent to one of skill in the art, other methods such as transient expression, which do not result in stable integration of the *BNM3* gene or which make direct use of the *BNM3* protein, may also be employed.

#### Use of *BNM3* Sequences to Target Gene Expression to the Embryo

Since *BNM3* genes are preferentially expressed in developing embryos (see example 3), a further application of this invention is the use of *BNM3* regulatory regions to target expression of at least one heterologous gene of interest to the developing embryo for any purpose, for example, but not limited to, altering embryo

- 33 -

and seed traits such as seed viability or size, composition of constituents of the seed, disease resistance, or the production of high value products such as vaccines antibodies, biopharmaceuticals or other specialty chemicals.

#### Use of *BNM3* Expression as a Marker for Embryogenic Cells

As shown in Examples 3 and 4, *BNM3* gene expression is detected during the earliest phase of plant embryogenesis and is itself sufficient to activate signal transduction cascades leading to embryo development. *BNM3* gene expression is therefore a specific marker for the entry of a plant cell into the embryogenic pathway.

*BNM3* expression is associated with embryo-forming cell divisions *in vitro* and *in vivo* and as such can be used to define culture conditions that alter the embryo-forming capacity of a tissue *in vitro*. Cells with embryogenic capacity or cells that undergo only a limited number of embryo-forming divisions are difficult to identify in the absence of structures that morphologically resemble embryos. However, these cells may be identified on the basis of *BNM3* expression. In this application, a vector containing the *BNM3* regulatory region, fused to a reporter gene, for example, but limited to, *GUS* (Jefferson *et al.*, 1987), *Luciferase* (Ow *et al.*, 1987) or *GFP* (Haselhoff and Amos, 1995) is transformed to a plant of interest. Homozygous transgenic lines exhibiting high levels of reporter gene expression in the embryo are cultured under *in vitro* conditions. Embryogenic cells, as well as culture conditions which facilitate or enhance the formation of embryogenic cells are identified on the basis of reporter gene expression within the cultured tissue.

A related application is the use of the *BNM3* gene as a marker in apomictic species for the identification of individual cells that are in the process of forming asexually-derived embryos. In this application, cells entering the autonomous embryo pathway are identified by mRNA *in situ* hybridization using a RNA probe derived from a *BNM3* gene sequence, by immunocytochemistry using an antibody directed against a *BNM3* protein, by transforming plants with a DNA construct containing a

- 34 -

gene fusion between *BNM3* regulatory regions and a reporter gene, or by any similar technique known to those skilled in the art.

#### Identification of Signal Transduction Components

Signal transduction components which activate or are activated by *BNM3* gene expression can be elucidated by identifying proteins and DNA sequences that interact with a *BNM3* gene and its protein product. These signal transduction components may be identified using techniques known to a person skilled in the art, including for example, but not limited to:

- yeast one hybrid screens for the isolation of proteins that bind to the *BNM3* regulatory regions to influence *BNM3* gene expression;
- genetic selection in yeast to identify genes that are direct targets of *BNM3* binding;
- DNA arrays or proteomics to identify genes which are activated in a *BNM3* signal transduction cascade; and
- yeast two hybrid screens to identify proteins that interact with *BNM3* to influence expression of downstream target genes.

Techniques for the analysis of the signal transduction components and signalling components are well known (see for example, Meijer *et al.* (1998), Lipshutz *et al.* (1999), and Anderson and Anderson (1998)).

Plants over-expressing the *BNM3* gene under control of a strong constitutive regulatory element such as, for example, but not limited to, the Cauliflower Mosaic Virus 35S promoter exhibit ectopic embryo formation, enhanced regeneration via organogenesis or a combination thereof (Examples 4 and 5). The ability of *BNM3* ectopic over-expression to induce both embryo formation and enhance regeneration processes can be used to identify mutants altered in their embryo-forming or regenerative capacity. In this application a vector construct consisting of a *BNM3* protein coding region under control of a regulatory element that is sufficient to promote either ectopic embryo formation or enhanced regeneration phenotype is made

- 35 -

and introduced into a plant of interest. Homozygous transgenic lines exhibiting a high penetrance of ectopic embryo formation, enhanced regeneration phenotype, or a combination thereof are identified. These lines are mutagenized by any available technique well known to the person skilled in the art, but which may include EMS mutagenesis, fast neutron mutagenesis, transposon mutagenesis or T-DNA mutagenesis. Mutagenized plants are then screened for alterations in the ectopic embryo formation or regeneration phenotype. These alterations include, for example, but not limited to, elimination or enhancement of the ability to promote ectopic asexual embryo formation or to regenerate in the absence of added growth regulators.

#### Heterologous Protein Expression System

Genetic control of the signal transduction pathway leading to embryogenesis and organogenesis in non-seed organs of transgenic plants may be activated by ectopic expression of a *BNM3* gene. Expression of a *BNM3* gene in association with a heterologous promoter can be used to produce altered seed components including for example, proteins, oils and other metabolites. Biotransformation of desired organs may also include altering the nutritive value of, for example leaves of forage crops, or it may be used to create alternative uses for crops. The use of promoters that are induced by the signal transduction cascade initiated by expression of *BNM3* can be used to express high-valued recombinant proteins in organs other than seeds. An example of one such promoter is the napin promoter, obtained from the 2S seed storage protein napin. The production of proteins initiated from a *BNM3*-induced cascade, may be achieved within organs exhibiting greater biomass than seeds. Therefore, this technology may be used to create alternatives for plants as crops.

Accordingly, the present invention further relates to a binary system in which the *BNM3* protein binds directly or indirectly to an embryo-expressed regulatory sequence (target sequence) and activate transcription of a chimeric gene construct in any plant cell, tissue or organ. Therefore, *BNM3* may be used to directly or indirectly activate transcription of a chimeric gene construct. This approach involves *BNM3*

- 36 -

interacting either directly with at least one target sequence from an embryo-expressed gene, or indirectly by initiating an embryogenic signal cascade that activates a transcription factor that in turn binds to and activates transcription from at least one target sequence. This binary system may be used for the expression of proteins in somatic tissues with the properties of expression in seeds.

In this application transgenic plants containing the *BNM3* gene under control of a constitutive regulatory element, for example, but not limited to the 35S promoter (35S:*BNM3*) are created to produce a *BNM3* activator line. *BNM3* expression may be demonstrated in a wide range of tissues in the *BNM3* activator lines by RNA gel blot analysis. Stable homozygous activator lines with high levels of *BNM3* expression are identified. Somatic tissues over-expressing *BNM3* may be examined for expression of other embryo-expressed genes, such as arabin (Guerche *et al.*, 1990), cruciferin (Pang *et al.*, 1988) or oleosin, or for morphological properties that are normally characteristic of seeds, such as the presence of lipid or protein bodies.

Transgenic plants of the same species to that used to generate the *BNM3* activator lines described above are also created which contain an embryo-expressed promoter fused to a gene of interest, to produce a gene of interest line. In order to help describe this embodiment, the gene of interest line expresses a reporter gene, such as *GUS*, and examples, which are not to be considered limiting, of such lines include ~~*Brassica napus* 2S albumin seed storage protein gene, *BngNAPI-GUS* fusion~~ (Baszczynski *et al.*, 1994) or a *SERK:GUS* fusion (Schmidt *et al.*, 1997; a non-seed expressed reporter construct such as *BNM1:GUS* (Treacy *et al.*, 1997) may be used as a negative control). The fidelity of expression of the gene of interest in the specific organs and tissues of these gene of interest lines is demonstrated for each construct. Stable homozygous lines with high levels of expression of the gene of interest expression are created.

Transgenic lines containing *BNM3* activator lines and gene of interest lines are crossed and the progeny seeds collected. *BNM3* gene expression, and in this example,



- 37 -

GUS activity, expression of other embryo-expressed genes, as well as the morphological characteristics of transformed tissues, are examined. *BNM3* expression in non-seed tissues typically activates both embryo development and expression of the gene of interest (e.g. GUS), however, activation of the expression of the gene of interest in the absence of morphologically discernible embryos may also be observed. Expression of the gene of interest, in the absence of morphologically discernible embryos provides initial evidence for direct interaction of *BNM3* with the target sequence.

Direct interaction of *BNM3* with a target sequence may also be demonstrated using transient expression of *BNM3* in plant protoplasts, along with the transient co-expression of an embryo-expressed promoter fused to a gene of interest (i.e. a gene of interest construct). *35S:BNM3* DNA and the gene of interest construct are introduced into protoplasts derived from non-seed cells, such as leaf mesophyll cells by electroporation. The expression of the gene of interest is examined after several hours to confirm activation of the target sequence. Direct interaction of *BNM3* with the target sequence may further be demonstrated by co-introducing the target sequence alone as competitor DNA.

In order to determine if tissues from different plant species may be transactivated by *BNM3*, *35S:BNM3* DNA and a reporter gene (for example, but not limited to *GUS*) construct may be introduced by microprojectile bombardment into somatic tissues of a plant. If *BNM3* interacts directly with a target sequence then expression of the reporter gene should coincide with transient expression of *BNM3* in all species and tissues.

Direct evidence for *BNM3*-target sequence interaction may also be obtained by isolation of *BNM3* protein expressed in bacteria, insect or yeast. *BNM3* is expressed in bacteria, insect, or yeast using commercially available expression systems and isolated to purity. Gel mobility shift assays (Gustavson *et al.*, 1991) are performed using a *BNM3*-target sequence, for example an embryo-expressed target

- 38 -

sequence, to demonstrate direct binding of BNM3 to the BNM3-target sequence. Footprint analyses may also be performed to locate the region of BNM3 binding. Fragments of target sequences that bind BNM3 may then be subcloned and used as competitors for BNM3 binding in transient assays described above.

The present invention will be further illustrated in the following examples. However it is to be understood that these examples are for illustrative purposes only, and should not be used to limit the scope of the present invention in any manner.

### Examples

#### General methods: *Microspore Embryo Culture*

*Brassica napus* c.v. Topas was used as the source of all plant material for microspore embryo culture. Donor plants for microspore culture were grown in a growth cabinet at 20 °C /15 °C (day/night) with a 16 h photoperiod (400 µE/m/s) provided by VHO cool white fluorescent lamps (165W, Sylvania) and incandescent bulbs (40W, Duro-test). Four weeks after germination the plants were transferred to growth cabinets under the same light conditions, but set at 10 °C /5 °C (day/night). Microspores and pollen were isolated and cultured as described in Keller *et al.* (1987), except that after 21 days in culture, cotyledon stage embryos were transferred to a maturation medium consisting of 1/2X NLN salts, 1% sucrose, 0.35 M mannitol and 5 µM ABA. Uninduced cultures (microspores and pollen continuing gametophytic development) and heat-stressed, non-embryogenic cultures (used for construction of the subtracted probe), were cultured from the same starting material as was used for the initiation of embryogenic cultures. Uninduced samples were obtained by culturing microspores and pollen for four days at 25 °C. Heat-stressed, non-embryogenic samples were obtained by culturing microspores and pollen for one day 25 °C, followed by three days 32 °C.

- 39 -

Samples of microspore and pollen cultured for less than 10 days were collected by centrifugation. Older samples containing globular, heart, torpedo and cotyledon stage microspore-derived embryos were collected by filtration through nylon meshes of various pore sizes as described in Ouellet *et al.* (1992). All other plant tissues were collected from greenhouse grown material. Seed material was obtained by hand pollinating flowers on the day of anthesis and collecting developing seeds on various days after pollination (DAP).

#### *Nucleic Acid Isolation and Analysis*

Total RNA was isolated using either a cesium chloride/guanidinium isothiocyanate procedure (Ouellet, 1992) or TRIZOL reagent (Gibco-BRL). RNA gel blot analysis was carried out by separation of 5 to 20 µg of total RNA per lane through 1.5% agarose gels containing 0.62 M formaldehyde, essentially as in Sambrook *et al.* (1989), followed by capillary transfer to Hybond-N nylon membranes (Amersham). Poly(A)<sup>+</sup> RNA was isolated from total RNA by oligo (dT)-cellulose chromatography (Sambrook, 1989).

Genomic DNA was isolated from leaf tissue as described in Fobert *et al.* (1991) and digested with the specified restriction enzymes using standard procedures (Sambrook, 1989). DNA gel blot analysis was carried out by electrophoresis of 10 µg DNA through 0.8% agarose gels followed by capillary transfer to Hybond-N membranes.

The partial 1.2 kb *BNM3A* cDNA insert was used as a probe for DNA and RNA gel blots. Hybridization to gel blots was carried out at 65 °C according to the Hybond-N protocol. The final wash conditions were 0.1X SSC, 65 °.

#### *Subtractive Probe Construction and cDNA Library Screening*

- 40 -

Poly (A) mRNA was isolated from late uninucleate microspores and early binucleate pollen that had been cultured for four days at 32 °C in order to induce embryogenesis (embryogenic sample) and used to synthesize first strand cDNA (Riboclone cDNA kit; Promega). The cDNA was then hybridized to a five-fold excess (by weight) of poly (A)<sup>+</sup> RNA from late uninucleate microspores and early binucleate pollen that had been cultured for one day at 25 °C, followed by three days at 32 °C to inactivate embryogenesis (non-embryogenic sample: Pechan *et al.*, 1991). The subtractive hybridization was performed essentially as described in Sambrook *et al.* (1989). The single-stranded cDNA recovered after subtraction was labelled with [ $\alpha$ -<sup>32</sup>P] dCTP using a random primers kit (BRL) and used as the subtracted probe for screening a Lambda phage cDNA library constructed from the same embryogenic sample described above (Boutillier, 1994). Triplicate nylon filter lifts (Hybond-N) from approximately  $1.5 \times 10^5$  plaque-forming units of the library were screened with the subtracted probe, with a random primers-labelled first strand non-embryogenic probe and with a random primers-labelled napin seed storage protein cDNA probe (pN2; Crouch, 1983). Napin mRNAs are prevalent in the embryogenic microspore library (Boutillier, 1994) and therefore plaques hybridizing to the napin probe were removed from the subsequent screening steps. Plaques hybridizing to the subtracted probe, but not to the non-embryogenic or napin probes, were selected and subjected to two subsequent rounds of differential screening using both the subtracted and non-embryogenic cDNA probes. DNA from selected Lambda clones was isolated (Sambrook, 1989), partially digested with *Eco* RI and *Xba* I and subcloned into pGEM-4Z (Promega).

Seven cDNAs comprising 6 unique genes, one of which comprised a truncated BNM3A cDNA, were identified. Two distinct, full length BNM3 cDNA clones (BNM3A and BNM3B) were subsequently obtained by stringent screening of circa  $2.5 \times 10^5$  plaque-forming units of a cDNA library (UniZAPII cDNA synthesis kit, Stratagene) constructed with mRNA from 10 day old globular to heart-stage microspore-derived embryos of *B. napus* c.v. Topas. The BNM3 cDNA inserts were rescued by *in vivo* excision into Bluescript SK(-) (Stratagene).

*Isolation of Genomic DNA sequences*

The Universal Genome Walker Kit (Clontech) was used to isolate genomic DNA fragments lying upstream of the *BNM3* ATG start codon. Pools of uncloned, adaptor-ligated *Brassica napus* cv Topas genomic DNA fragments were constructed and used to isolate *BNM3* genomic sequences by nested PCR. The primary PCR made use of the outer adaptor primer (AP1) supplied by the manufacturer and a *BNM3* specific primer with the sequence:

5'-GAGGCAGCGTCGGATCGTAACAGTACTCT-3' (SEQ ID NO:6).

The nested PCR made use of the nested adaptor primer (AP2) supplied by the manufacturer and a *BNM3* specific primer with the sequence:

5'-CATAAGGAGAGAGAGAAAAGCCTAACCACT - 3' (SEQ ID NO:7).

The primary PCR mixture was then diluted 1:50 and used as template for nested PCR. Both the primary and nested PCRs were performed as recommended by the manufacturer. The nested PCR products were cloned into the pGEMT-Easy vector (Promega) and sequenced. PCR products corresponding to the 5' untranslated genomic regions of both *BNM3A* and *BNM3B* cDNAs were identified.

The genomic DNA sequence spanning the *BNM3A* ATG translational start and TAG translational stop codons was isolated by PCR from *B. napus* cv Topas genomic DNA using *Pfu* polymerase (Stratagene) and the following primer combination:

5'-ACCAAGAAGTCGTTAGATC-3' (SEQ ID NO:8); and

5'-AAGCATATAACTAAAGATC-3' (SEQ ID NO:9).

The primers were used under standard PCR conditions. The PCR products were cloned into the pGEMT-Easy vector and sequenced.

*Plasmid Construction for Plant Transformation*

The construction of a plasmid vectors containing the *BNM3* cDNAs under control of either a *POLYUBIQUITIN* or Cauliflower Mosaic virus 35S promoter are

- 42 -

described below. The plasmid pRAP2TUBI contains a modified *Helianthus annuus* *POLYUBIQUITIN* promoter (Binet *et al.*, 1991) in the plasmid pRAP2T. The plasmid pRAP2T consists of the pUCAP plasmid (van Engelen *et al.*, 1995) and a nopaline synthase (nos) terminator inserted into the *Sac* I and *Eco* RI restriction sites. A PCR fragment of the *POLYUBIQUITIN UbbI* promoter comprising the 5' end of the promoter to 7 bp from the 3' end of the first exon was amplified from the vector using an M13 reverse primer and the UBIQ-3' primer:

5'-CCATGGATCCAGAGACGAAGCGAAAC-3' (SEQ ID NO:10)

which includes introduced *Nco* I and *Bam* HI restriction sites. The *POLYUBIQUITIN* promoter fragment was digested with *Pst* I and *Bam* HI, gel purified and ligated into the *Pst* I and *Bam* HI sites of pRAP2T, creating the vector pRAP2TUBIH. The full-length *BNM3B* cDNA was digested with *Eco* RI and *Xho* I restriction enzymes, blunted with Klenow enzyme, gel purified and ligated into the *Sma* I site of pRAP2TUBI making the plasmid pKB1S. An *Asc* I/*Pac* I DNA restriction fragment containing the modified *POLYUBIQUITIN* promoter, the *BNM3B* cDNA and the nos terminator was gel purified, and ligated to the *Asc* I/*Pac* I digested binary vector pBINPLUS (van Engelen *et al.*, 1995), creating the plasmid pKBBIN1S.

The construction of a vector containing the *BNM3A* cDNA under control of a double enhanced 35S promoter and AMV translational enhancer was as follows. A *Hind* III/*Xba* I DNA restriction fragment containing the double 35S promoter and the AMV translational enhancer from plasmid pBJ525 (Datla *et al.*, 1993) was ligated to *Hind* III/*Xba* I digested pRAP2T, creating the plasmid pRAP2T35S. An *Nco* I site was introduced into the *BNM3A* cDNA clone by site directed mutagenesis. The sequence of the BNM3ANCO1 primer used for mutagenesis is:

5'-ACTCCATGGATAATAACTGGTTAGGC-3' (SEQ ID NO:11).

A second primer, BNM3AHINDIII:

5' - AAATTCTCAAGCTTTGGTCCATCTTG-3' (SEQ ID NO12)

was used together with the BNM3ANCO1 primer to amplify a 305 bp fragment of the *BNM3A* cDNA. This PCR fragment was digested with *Nco* I and *Hind* III and ligated to *Nco* I/*Kpn* I cut pRAP2T35S and a *Hind* III/*Kpn* I fragment containing the region of

- 43 -

the *BNM3A* cDNA downstream of the *Hind* III site, creating the vector p35S:BNM3. p35S:BNM3 was digested *Asc* I and *Pac* I restriction enzymes and the fragment containing the double 35S promoter, the AMV translational enhancer, the *BNM3A* cDNA and the nos terminator was gel purified and ligated to the *Asc* I/*Pac* I digested binary vector pBINPLUS, creating the plasmid p35S:BNM3BIN.

Both the pKBBIN1S and p35S:BNM3BIN plasmids were transferred to *Agrobacterium tumefaciens* C58C1 strain carrying the disarmed Ti plasmid pMP90 and used in transformation experiments.

#### *Plant Transformation*

*Arabidopsis thaliana* ecotype C24 was used as the recipient in transformation experiments. Plants were transformed using either the floral dip method described in Clough and Bent (1998) or the root transformation method described in Vergunst *et al.* (1998).

Transgenic *Brassica napus* c.v. "Topas" plants were produced by *Agrobacterium tumefaciens*-mediated transformation of microspore-derived embryos. Microspore-derived embryos were cultured for 5 weeks at a density of approximately 1000 embryos per ml. Overnight cultures of *Agrobacterium* were diluted 100 times in B5 medium containing 9% sucrose. Embryos were co-cultivated with the diluted bacteria for 48 hours at 24°C in darkness, with slow shaking. The embryos were then transferred to NLN13 medium supplemented with 350 mg/L cefotaxim and 200 mg/L vancomycin for at least two weeks in darkness at 25°C.

Embryos were germinated in weak light at 25°C for about 2 weeks on solid B5 medium supplemented with 2% sucrose, cefotaxim (200 mg/L) and vancomycin (100 mg/L). Well developed hypocotyls from germinated embryos were isolated and transferred to fresh germination medium supplemented with 100 mg/L kanamycin. After two weeks on this medium, explants were subcultured to a similar medium

- 44 -

supplemented with kanamycin (25 mg/L). Green, putative transgenic, secondary embryos become visible after one month of selection.

### *Microscopy*

All plant material was fixed overnight at 4 °C in 0.1 M phosphate buffer pH 7.0 containing 4% paraformaldehyde. Samples were washed in 0.1 M phosphate buffer and then dehydrated in a graded ethanol series to 100% ethanol. Samples for scanning electron microscopy were critical point dried in liquid CO<sub>2</sub> (Balzers CPD020), and mounted on SEM stubs using conductive carbon glue. Samples were coated with 30 nm palladium/gold using a Polaron E5100 sputter coater. Samples were observed in a JEOL JSM 5200 scanning electron microscope with an acceleration voltage of 15 kV. Digital images were obtained using Orion Framegrabber. Samples for light microscopy were embedded in Technovit 7100 (Kulzer). Sections were stained for 10 seconds in 1% Toluidine blue in 1% sodiumtetraborate, rinsed with water and mounted in Euparal. Digital images were recorded using a Sony 3 CCD camera.

### *Regeneration Experiments*

Wild-type and transgenic arabidopsis seeds were surface sterilized, plated on 1/2 MS media containing 20% sucrose (1/2MS-20) and grown at 21 °C with the plates inclined at a 60° angle. Eight wild-type seedlings and eight seedlings from each of seven independent transgenic lines were harvested 10 days after germination and separated into root, hypocotyl and leaf explants. This material was then divided into two batches. Half of the explants were continuously cultured on B5 media containing 20% glucose (B5-20). Explants were transferred to fresh B5-20 media every two weeks. The remaining explants were cultured on B5-20 containing plant growth regulators in order to induce shoot regeneration (Vergunst *et al.*, 1998). These explants were first placed on callus inducing media (CIM; high auxin to cytokinin ratio) for two days and then transferred to shoot inducing media (SIM; high cytokinin



- 45 -

to auxin ratio) for the remainder of the culture period. Explants were transferred to fresh SIM media every two weeks.

*Example 1: Isolation and Characterization of the BNM3 Genes from Brassica napus*

A subtractive screening approach was used to isolate genes preferentially expressed during the induction of *Brassica napus* c.v. Topas microspore embryogenesis (Figure 1). Two types of microspore cultures were used in the construction of a subtracted probe: embryogenic and non-embryogenic. Embryogenic cultures were obtained by subjecting late uninucleate microspores and early binucleate pollen to a 4 day, 32 °C heat stress treatment. The non-embryogenic sample was obtained by culturing the same starting population of late uninucleate microspores and early binucleate pollen for 1 day at 25 °C followed by 3 days at 32 °C (Pechan *et al.*, 1991). Poly(A) mRNA was isolated from the embryogenic sample and used to synthesize first strand cDNA. The cDNA was then hybridized to an excess of poly(A)<sup>+</sup> RNA isolated from a non-embryogenic microspore/pollen sample. The non-hybridizing, single stranded cDNA, enriched for sequences present in the embryogenic sample, but absent or present at a much lower level in the non-embryogenic sample, was recovered, radioactively labelled and used as a subtracted probe for screening a cDNA library derived from the embryogenic sample described above. ~~Plaques hybridizing to the subtracted probe, but not to a probe derived from~~ the non-embryogenic sample, were selected and subjected to two subsequent rounds of differential screening. Seven independent cDNA clones, comprising six unique DNA sequences were found to be differentially expressed between the embryogenic and non-embryogenic samples. One of these clones, 42A1, later renamed *BNM3A* (for *Brassica napus* microspore embryo), was further characterized.

*Example 2: The BNM3 genes encode new members of the AP2 domain class of transcriptional activators*

- 46 -

A single *BNM3* cDNA clone, *BNM3A*, was isolated after screening an embryogenic microspore cDNA library with a subtracted probe enriched for genes expressed in embryogenic microspores and pollen. The discrepancy between the size of the cDNA clone (1.2 kb) and the size of the transcript detected on RNA gel blots (2.2 kb) indicated that this clone did not represent a full-length cDNA. Two longer cDNA clones, corresponding to the full length cDNA of the clone originally isolated, *BNM3A* (SEQ ID NO. 1), and a new clone, *BNM3B* (SEQ ID NO. 3), were isolated from a 10 day old *Brassica napus* microspore embryo cDNA library. The alignment of the DNA sequence of these clones is shown in Figure 2. The two *BNM3* cDNA clones are 2011 and 1992 nt in length, and are 97% similar at the nucleotide level, differing only slightly in the length and sequence of their 5' and 3' untranslated regions. Both cDNAs potentially encode 579 amino acid polypeptides (predicted molecular mass of 63.9 kDa, pI of 5.7) that are 97% similar at the amino acid level (Figure 3).

The genomic complexity of the *BNM3* genes was determined by hybridization of the *BNM3* cDNAs to gel blots containing *B. napus* genomic DNA (Figure 4). The *BNM3* cDNAs hybridize to two DNA fragments under high stringency conditions. The two hybridizing fragments represent the two *BNM3* genes, *BNM3A* and *BNM3B*. *B. napus* is an amphidiploid species derived from the hybridization of the diploid *B. rapa* and *B. oleracea* genomes, thus the two *BNM3* sequences are likely derived from a single copy locus in each of the parental diploid progenitors.

Search of the sequence databases indicated that the *BNM3* translation products contain two copies of an AP2 domain (Figure 3). The AP2 domain was first identified in APETALA2 (AP2), an arabidopsis protein that regulates meristem identity, floral organ specification, seedcoat development and floral homeotic gene expression (Jofuku *et al.*, 1994; WO 98/07842), and has since been identified in a wide range of proteins with diverse functions. These functions range from the activation of genes involved in stress (Zhou, 1997; Stockinger, 1997) and ethylene response (Ohme-Takagi, 1995) to the regulation of leaf, floral and ovule development (Moose, 1996; Jofuku, 1994; Elliot, 1996; Klucher, 1996). The AP2 domain is a 56-68 amino acid

- 47 -

repeated motif containing at least two conserved regions: a highly basic YRG element, containing a conserved YRG amino acid motif and the RAYD element. The RAYD element contains a conserved central core of 18 amino acids that is predicted to form an amphipathic  $\alpha$ -helix, a structure that is thought to mediate protein-protein interactions. The ability of a number of AP2 domain containing proteins to bind DNA, coupled with the presence of putative nuclear localization signals and acidic regions that may function as transcriptional activators suggests these proteins function as transcription factors.

Two phylogenetically distinct classes of AP2 domain proteins, consisting of either one AP2 domain (EREBP-like) or two AP2 domains connected by a linker region (AP2-like), have been identified (Zhou, 1997). BNM3 belongs to the latter class. Search of the databases with the region corresponding to the two AP2 domains and linker region of BNM3 reveals that BNM3 is most similar to the arabidopsis AINTEGUMENTA (ANT; Elliot, 1996; Klucher, 1996) and the *Zea mays* ZMMHCF1 AP2 domain containing protein. (ZM; Daniell, 1996) Figure 5 shows an alignment of the two AP2 domains of BNM3 with those of other proteins that contain two AP2 domains. BNM3 shares 85% amino acid sequence similarity with ANT and 88% with ZMMHCF1 in this region, but only 66% amino acid similarity with AP2 and GLOSSY15 in this region. A 10 amino acid insertion in the first AP2 domain of the and BNM3 proteins further distinguishes these three proteins from other AP2 domain containing proteins (Elliot, 1996). The BNM3, AINTEGUMENTA and ZMMHCF1 proteins also share a small hydrophobic amino acid motif, LG/SFSL, in their amino terminal regions, but otherwise show no significant similarity in their DNA or amino acid sequences outside of the AP2 domains and linker. These results indicate that the BNM3 sequences encode unique members of the AP2 domain family of proteins.

A pairwise alignment of BNM3B cDNA and amino acid, sequences with ANT or ZMMHCF-1 sequences indicated that for the BNM3B nucleotide sequence:

- 48 -

- there is a 56% identity with ANT cDNA (over the 1905 nucleotides of ANT) and a 58% identity with ZMMHCF1 cDNA (over the 1773 nucleotide sequence of ZM);

and for the BNM3B amino acid sequence:

- there is a 41% identity of the BNM3B protein with ANT protein (over the 555 amino acid sequence of ANT), and a 46% identity with ZMMHCF1 protein (over 485 amino acid sequence of ZM).

*Example 3: The BNM3 genes are preferentially expressed in developing embryos*

RNA gel blot analysis (Figure 6) was used to determine the pattern of *BNM3* gene expression during microspore-derived embryo development, seed development, and in non-seed tissues. Both analyses indicate that the *BNM3* genes are preferentially expressed in developing embryos.

RNA gel blot analysis indicates that *BNM3* mRNAs are detected in microspore cultures induced to undergo embryogenesis, as well as in the subsequent globular, heart, torpedo and cotyledon stages of microspore-derived embryo development (Figure 6A). *BNM3* mRNAs are not detected in non-embryogenic microspore cultures, in freshly isolated microspores and pollen, or in microspores and pollen continuing gametophytic development in culture (Figure 6A). RNA gel blot analysis of developing seeds shows that *BNM3* expression is first detected 14 days after pollination (14 DAP), corresponding to the heart stage of embryo development. *BNM3* expression increases during the early (21 DAP) and mid-cotyledon (28 DAP) stages of embryo development and remains constant thereafter (Figure 6B). *BNM3* transcripts were not detected in any of the non-seed tissues tested, reflecting the low level or absence of transcripts in these tissues.

*Example 4: Expression of BNM3 in Vegetative Tissues Promotes Asexual Embryo Formation*

- 49 -

In order to determine the function the *Brassica napus* *BNM3* proteins, the *BNM3* cDNAs were placed under the control of two separate constitutive promoter constructs, a modified sunflower *POLYUBIQUITIN* promoter construct (hereafter referred to as *UBI:BNM3*) and a double enhanced 35S promoter construct containing an AMV translational enhancer (hereafter referred to as *35S:BNM3*), and introduced into arabidopsis. Analysis of the phenotype of the transformants indicates that ectopic over expression of the *BNM3* cDNAs promotes the formation of somatic embryos on vegetative structures such as cotyledons, petioles, leaf blades and the shoot apical meristem (Figure 7). The frequency of transformants producing ectopic embryos, as well as the penetrance of the ectopic embryo phenotype, was greater when the *BNM3* gene was expressed under control of the stronger double enhanced 35S promoter-AMV translational enhancer, as compared to the *POLYUBIQUITIN* promoter. Thus a high threshold level of protein product is required to increase the frequency and penetrance of the ectopic embryo phenotype.

*BNM3*-derived ectopic embryos contain all of the organ systems and tissue layers found in the developing zygotic embryo. *BNM3*-derived ectopic embryos are bipolar (Figures 7D and E) and consist of an axis, comprised of the hypocotyl and radicle regions, shoot and root meristems, and cotyledons (Figure 7E). In addition, each organ system contains the characteristic radial arrangement of three specialized tissue layers (epidermis, ground parenchyma and provascular tissue) found in zygotic embryos (Figure 7E). Continued expression of the *BNM3* gene within the developing ectopic embryo leads to a reiteration of the embryo-forming process, with the result that new embryos are continuously formed on the surface of pre-existing embryos (Figure 7D and E). These results provide conclusive evidence that expression of a single gene, *BNM3*, is sufficient to initiate a signal transduction cascade leading to the formation of fully differentiated asexually-derived embryos.

*Example 5: Expression of BNM3 Increases the Regeneration Capacity of Plant Tissues*

- 50 -

We examined the effect of *BNM3* gene expression on the ability of arabidopsis plants to regenerate shoots *in vitro* in the presence or absence of added growth regulators. Leaf, root and hypocotyl explants from 10 day old seedlings of wild-type arabidopsis and transgenic arabidopsis lines expressing *BNM3* under control of the *POLYUBIQUITIN* promoter were placed on media containing growth regulators to induce first callus formation and then shoot organogenesis. Root explants from transgenic lines show at least a 5-fold increase in shoot regeneration in the presence of hormones as compared to wild-type root explants. (Figure 8A). These shoots also developed faster in the transgenic explants as compared to the wild-type. Wild-type leaf and hypocotyl explants responded by producing callus on the cut end of the petiole (Figure 8B). In contrast, explants from transgenic lines immediately produced new shoots (Figure 8B) or roots from the cut end of the petiole. Transgenic explants that initially produced roots eventually also produced shoots.

Transgenic explants were also able to regenerate in absence of added growth regulators. Wild-type leaf and hypocotyl explants placed on medium lacking growth regulators occasionally produced callus or roots at the cut end of the leaf petiole, however shoots did not regenerate from these structures (Figure 8C,D). Wild-type roots greened and formed thickened nodule-like structures at the junction with lateral roots, but did not develop further. In contrast, transgenic explants placed on media lacking growth regulators regenerated shoots either from the cut end of the leaf and hypocotyl explants or from the nodule-like structures of root explants (Figure 8C,D).

The present invention has been described with regard to preferred embodiments. However, it will be obvious to persons skilled in the art that a number of variations and modifications can be made without departing from the scope of the invention as described herein.

## References

- 51 -

Anderson, N.L. and Anderson, N.G. (1998). Proteome and proteomics: new technologies, new concepts and new words. *Electrophoresis* 19, 1853-1861.

Angenent, G.C., Franken, J., Busscher, M., van Dijken, A., van Went, J.L., Dons, H.J.M. and van Tunen, A.J. (1995). A novel class of MADS box genes involved in ovule development in *Petunia*. *Plant Cell* 7, 1569-1582.

Aoyama, T. and Chua, N.H. (1997). A glucocorticoid-mediated transcriptional induction system in transgenic plants. *Plant J.* 2, 397-404.

Baszczynski, C.L., and Fallis, L. (1990). Isolation and nucleotide sequence of a genomic clone encoding a new *Brassica napus* napin gene. *Plant Mol. Biol.* 14, 633-635.

Binet, M.N., Lepetit, M., Weil, J.H. and Tessier, L.H. (1991). Analysis of a sunflower polyubiquitin promoter by transient expression. *Plant Sci.* 79, 87-94.

Boutilier, K.A., Gines, M.J., Demoor, J.M., Huang, B., Baszczynski, C.L., Iyer, V.N. and Miki, B.L. (1994). Expression of the BnmNAP subfamily of napin genes coincides with the induction of *Brassica* microspore embryogenesis. *Plant Mol. Biol.* 26, 1711-1723.

---

Brandstatter, I. and Kieber, J.J. (1998). Two genes with similarity to bacterial response regulators are rapidly and specifically induced by cytokinin in *Arabidopsis*. *Plant Cell* 10, 1009-1019.

Caddick, M.X., Greenland, A.J., Jepson, I., Krause, K.P., Qu, N., Riddell, K.V., Salter, M.G., Schuch, W., Sonnewald, U., and Tomsett, A.B. (1998). An ethanol inducible gene switch for plants used to manipulate carbon metabolism. *Nature Biotech.* 16, 177-180.

- 52 -

Chaudury, A.M., Letham, D.S., Craig, S. and Dennis, E.S. (1993). amp-1-a mutant with higher cytokinin levels and altered embryonic pattern, faster vegetative growth, constitutive photomorphogenesis and precocious flowering. *Plant J.* 4, 907-916.

Clough, S.J. and Bent, A.F. (1998). Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J.* 16, 735-743.

Custers, J.B.M., Oldenhof, M.T., Schrauwen, J.A.M., Cordewener, J.H.G., Willems, G.J. and van Lookeren Campagne, M.M. (1997) Analysis of microspore-specific promoters in tobacco. *Plant Mol. Biol* 35, 689-699.

Crouch, M.L., Tenbarge, K.M., Simon, A.E. and Ferl, R. (1983). cDNA clones for *Brassica napus* seed storage proteins: evidence from nucleotide sequence analysis that both subunits of napin are cleaved from a precursor polypeptide. *J. Mol. Appl. Genet.* 2, 273-283.

Daley, M., Knauf, V.C., Summerfelt, K.R. and Turner, J.C. (1998). Co-transformation with one *Agrobacterium tumefaciens* strain containing two binary plasmids as a method for producing marker free transgenic plants. *Plant Cell Rep.* 17, 489-496.

Daniell, T.J., Fordham-Skelton, A.P., Vergani, P. and Edwards, R. (1996). Isolation of a maize cDNA (accession no. Z47554) (PGR 96-013) encoding APETALA-2-like binding domains by complementation cloning of an L-isoaspartyl methyltransferase-deficient mutant of *Escherichia coli*. *Plant Phys.* 110, 1435.

Datla, R.S.S., Bekkaoui, F., Hammerlindl, J.K., Pilate, G., Dunstan, D.I. and Crosby, W.L. (1993). Improved high-level constitutive foreign gene expression in plants using an AMV RNA4 untranslated leader sequence. *Plant Sci.* 94, 139-149.



- 53 -

DeBlock, M. DeBrower, D. and Tenning, P. (1989). Transformation of *Brassica napus* and *Brassica oleracea* using *Agrobacterium tumefaciens* and the expression of the bar and neo genes in the transgenic plants. *Plant Physiol.* 91: 694-701.

Dellaert, L.M.W. (1981) Comparison of X-ray and fast neutron-induced mutant spectra. Experiments in *Arabidopsis thaliana* (L.) Heynh. *Arabidopsis Inf. Ser.*, 18, 16-36.

Doan, D. N. P., Linnestad, C., and Olsen, O.-A. (1996). Isolation of molecular markers from the barley endosperm coenocyte and the surrounding nucellus cell layers. *Plant Mol. Biol.* 31: 877-886.

Elliot, R.C., Betzner, A.S., Huttner, E., Oakes, M.P., Tucker, W.Q.J., Gerentes, D., Perez, P. and Smyth, D.R. (1996). AINTEGUMENTA, an APETALA2-like gene of *Arabidopsis* with pleiotropic roles in ovule development and floral organ growth. *Plant Cell* 8, 155-168.

Federoff, N., Furtek, D., and Nelson O. (1984). Cloning of the bronze locus in maize by a simple and general procedure using the transposable controlling element Ac. *Proc. Natl. Acad. Sci. USA* 81, 3825-3829.

---

Feldman, K.A., Marks, M.D., Christianson, M.L., and Quatrano, R.S. (1989). A dwarf mutant *Arabidopsis* generated by T-DNA insertion mutagenesis. *Science* 243, 1351-1354.

---

Fobert, P.R., Miki, B.L., and Iyer, V.N. (1991). Detection of gene regulatory signals in plants revealed by T-DNA-mediated fusions. *Plant Mol. Biol.* 17, 837-851.

Gatz, C. (1997). Chemical control of gene expression. *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 48, 89-108.

- 54 -

Gatz, C. and Lenk, I.R.P. (1998). Promoters that respond to chemical inducers. Trends Plant Sci. 3, 352-358.

Guerche, P., Tire, C., Grossi De Sa, F., De Clercq, A., Van Montagu, M. and Krebbers, E. (1990). Differential expression of the Arabidopsis 2S albumin genes and the effect of increasing gene family size. Plant Cell 2, 469-478.

Gustavsson, H.O., Ellerstrom, M., Stulberg, K., Ezcurra, I., Koman, A., Hoglund, A., Rask, L. and Josefsson, L.-G. (1991). Distinct sequence elements in a napin promoter interact in vitro with DNA-binding proteins from Brassica napus. Physiol. Plant 82, 205-212.

Haseloff, J. and Amos, B. (1995). GFP in plants. Trends Genet. 11, 328-329.

Jefferson, R.A. and Bicknell, R. (1996). The potential impacts of apomixis: a molecular genetics approach. In The Impact of Plant Molecular Genetics, B.W.S. Sobral, ed (Boston: Birkhauser), pp. 87-101.

Jefferson, R.A., Kavanagh, T.A. and Bevan, M.W. (1987). GUS fusions:  $\beta$ -glucuronidase as a sensitive and versatile gene fusion marker in higher plants. EMBO J. 6, 3901-3907.

Jofuku, K.D., den Boer, B.G.W., van Montagu, M. and Okamoto, J.K. (1994). Control of Arabidopsis flower and seed development by the homeotic gene APETALA2. Plant Cell 6, 1211-1225.

Kakimoto, T. (1996). CKI7, a histidine kinase homolog implicated in cytokinin signal transduction. Science 274, 982-985.

- 55 -

Keller, W.A., Fan, Z., Pechan, P., Long, N., Grainger, J. (1987). An efficient method for culture of isolated microspores of *Brassica napus*. Proceedings of the 7th International Rapeseed Congress. Poznan, Poland. Vol. 1, 152-157.

Klein, T.M., Wolf, E.D., Wu, R. and Sanford, J.C. (1987). High-velocity microprojectiles for delivering nucleic acids into living cells. *Nature* 327,70-73

Klimyuk, V.I. and Jones, J.D.G. (1997). AtDMC1, the Arabidopsis homologue of the yeast DMC1 gene: characterization, transposon-induced allelic variation and meiosis-associated expression. *Plant J.* 11, 1-14.

Klucher, K.M., Chow, H., Reiser, L. and Fischer, R.L. (1996). The AINTEGUMENTA gene of Arabidopsis required for ovule and female gametophyte development is related to the floral homeotic gene APETALA2. *Plant Cell* 8, 137-153.

Koltunow, A.M., Bicknell, R.A. and Chaudhury, A.M. (1995). Apomixis: molecular strategies for the generation of genetically identical seeds without fertilization. *Plant Physiol.* 108,1345-1352.

Koltunow, A.M. (1993). Apomixis: embryo sacs and embryos formed without meiosis or fertilization in ovules. *Plant Cell* 5, 1425-1437.

Korneef, M., Hanhart, C.J. and Thiel, F. (1989). A genetic and phenotypic description of eceriferum (cer) mutants in *Arabidopsis thaliana*. *J. Hered.* 80, 118-122.

Lightner J., and Caspar, T. (1988) Seed Mutagenesis of Arabidopsis. In Arabidopsis Protocols, J.M. Martinez-Zapater and J. Salinas eds (Totowa, USA: Humana Press).

Lipshultz, R.J., Fodor, S.P.A., Gingeras, T.R. and Lockhart, D.J. (1999). High density synthetic oligonucleotide arrays. *Nature Genetics* 21, 20-24.

- 56 -

Lotan, T., Ohto, M., Matsudaira Yee, K., West, M.A.L., Lo, R., Kwong, R.W., Yamagishi, K., Fischer, R.L., Goldberg, R.B. and Harada, J.J. (1998). Arabidopsis LEAFY COTYLEDON1 is sufficient to induce embryo development in vegetative cells. *Cell* 93, 1195-1205.

Meijer, A.H., Ouwkerk, P.B.F. and Hoge, J.H.C. (1998). Vectors for transcription factor cloning and target site identification by means of genetic selection in yeast. *Yeast* 14, 1407-1415.

Moose, S.P. and Sisco, P.H. (1996). Glossy15, an APETALA2-like gene from maize that regulates leaf epidermal cell identity. *Genes Dev.* 10, 3018-3027.

Mordhorst, A.P., Toonen, M.A.J. and de Vries, S.C. (1997). Plant embryogenesis. *Crit. Rev. Plant Sci.* 16, 535-576.

Mordhorst, A.P., Voerman, K.J., Hartog, M.V., Meijer, E.A., van Went, J., Koomneef, M., and de Vries, S.C. (1998). Somatic embryogenesis in *Arabidopsis thaliana* is facilitated by mutations in genes repressing meristematic cell divisions. *Genetics* 149, 549-563.

Ogas, J., Cheng, J.-C., Sung, R.Z. and Somerville, C. (1997). Cellular differentiation regulated by gibberellin in the *Arabidopsis thaliana* pickle mutant. *Science* 277, 91-94.

Ohme-Takagi, M. and Shinshi, H. (1995). Ethylene-inducible DNA binding proteins that interact with an ethylene responsive element. *Plant Cell* 7, 173-182.

Oldenhof, M.T., de Groot, P.F.M., Visser, J.H., Schrauwen, J.A.M. and Wullems, G.J. (1996). Isolation and characterization of a microspore-specific gene from tobacco. *Plant Mol. Biol.* 31, 213-225.

- 57 -

Ouellet, T., Rutledge, R.G. and Miki, B.L. (1992). Members of the acetohydroxyacid synthase multigene family of *Brassica napus* have divergent patterns of expression. *Plant J.* 2, 321-330.

Ow, D.W., Jacobs, J.D. and Howell, S.H. (1987). Functional regions of the cauliflower mosaic virus 35S RNA promoter determined by use of the firefly luciferase gene as a reporter for promoter activity. *Proc. Natl. Acad. Sci. USA* 84, 4870-4874.

Ozias-Akins, P., Lubbers, E.L., Hanna, W.W. and McNay, J.W. (1993). Transmission of the apomictic mode of reproduction in *Pennisetum*: co-inheritance of the trait and molecular markers. *Theor. Appl. Genet.* 85, 632-638.

Pang, P., Pruitt, R. and Meyerowitz, E. (1988). Molecular cloning, genomic organisation, expression and evolution of the 12S storage protein genes of *Arabidopsis thaliana*. *Plant Mol. Biol.* 11, 805-820.

Peacock, W.J., Ming, L., Craig, S., Dennis, E., Chaudury, A.M. (1995). A mutagenesis programme for apomixis genes in *Arabidopsis*. In *Proceedings Symposium on Induced Mutations and Molecular Techniques for Crop Improvement*, (Vienna: International Atomic Energy Agency), pp. 117-125

---

Pechan, P.M., Bartels, D., Brown, D.C.W. and Schell, J. (1991). Messenger-RNA and protein changes associated with induction of *Brassica* microspore embryogenesis. *Planta* 184, 161-165.

Roberts, M.R., Foster, G.D., Blundell, R.P., Robinson, S.W., Kumar, A., Draper, J. and Scott, R. (1993). Gametophytic and sporophytic expression of an anther-specific *Arabidopsis thaliana* gene. *Plant J.* 3, 111-120.

- Rounsley, S.D., Ditta, G.S. and Yanofsky, M.F. (1995). Diverse roles for MADS box genes in *Arabidopsis* development. *Plant Cell* 7, 1259-1269.
- Salter, M.G., Paine, J.A., Riddell, K.V., Jepson, I., Greenland, A.J., Caddick, M.X., Tomsett, A.B. (1998). Characterisation of the ethanol-inducible alc gene expression system for transgenic plants. *Plant Journal* 16, 127-132.
- Sambrook, J., Fritsch, E.F. and Maniatis, T. (1989). *Molecular Cloning: A Laboratory Manual*, second edition. (Cold Spring Harbor: Cold Spring Harbor Laboratory Press).
- Schmidt, E.D.L., Guzzo, F., Toonen, M.A.J. and de Vries, S.C. (1997). A leucine-rich repeat containing receptor-like kinase marks somatic plant cells competent to form embryos. *Development* 124, 2049-2062.
- Stockinger, E.J., Gilmour, S.J. and Thomashow, M.F. (1997). *Arabidopsis thaliana* CBF1 encodes an AP2 domain-containing transcriptional activator that binds to the C-repeat/DRE, a cis-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit. *Proc. Natl. Acad. Sci. USA* 94, 1035-1040.
- Taylor, R.L. (1967). The foliar embryos of *Malaxis paludosa*. *Can. J. Bot.* 45, 1553-1556.
- Treacy, B.K., Hattori, J., Prud'homme, I., Barbour, E., Boutilier, K., Baszczynski, C.L., Huang, B., Johnson, D.A. and Miki, B.L. (1997). Bnm1, a Brassica pollen-specific gene. *Plant Mol. Biol.* 34, 603-611.
- Twell, D., Wing, R., Yamaguchi, J. and McCormick, S. (1989). Isolation and expression of an anther-specific gene from tomato. *Mol. Gen. Genet.* 217, 240-245.

- 59 -

- Ulmasov, T., Murfett, J., Hagen, G. and Guilfoyle, T. (1997). Aux/IAA proteins repress expression of reporter genes containing natural and highly active synthetic auxin response elements. *Plant Cell* 9, 1963-1971.
- van Engelen, F.A., Molthoff, J.W., Conner, A.J., Nap, J.P., Pereira, A. and Stiekema, W.J. (1995). pBINPLUS: an improved plant transformation vector based on pBIN19. *Transgenic Res.* 4, 288-290.
- Vergunst, A.C., de Waal, E.C. and Hooykaas, P.J.J. (1998). Root transformation by *Agrobacterium tumefaciens*. In *Arabidopsis Protocols*, J.M. Martinez-Zapater and J. Salinas, eds (Totowa, USA: Humana Press).
- Xu, D., McElroy, D., Thornburg, R.W. and Wu, R.C.S. (1993). Systemic induction of a potato pin2 promoter by wounding, methyl jasmonate, and abscisic acid in transgenic rice plants. *Plant Mol. Biol.* 22, 573-588.
- Xu, H., Knox, R.B., Taylor, P.E. and Singh, M.B. (1995). Bcp1, a gene required for male fertility in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* 92, 2106-2110.
- Yarbrough, J.A. (1932). Anatomical and developmental studies of the foliar embryos of *Bryophyllum calycinum*. *Amer. J. Bot.* 19, 443-453.
- 
- Yeung, E.C. (1995). Structural and developmental patterns in somatic embryogenesis. In *In Vitro Embryogenesis in Plants*, T.A. Thorpe, ed (Dordrecht: Kluwer Academic Publishers), pp. 205-247.
- Zarsky, V., Garrido, D., Eller, N., Tupy, J., Vicente, O., Schaffl, F. and Heberle-Bors, E. (1995). The expression of a small heat shock gene is activated during induction of tobacco pollen embryogenesis by starvation. *Plant Cell Environ.* 18, 139-147.

- 60 -

Zhou, J., Tang, X. and Martin, G.B. (1997). The Pto kinase conferring resistance to tomato bacterial speck disease interacts with proteins that bind a cis-element of pathogenesis-related genes. *EMBO J.* 16, 3207-3218.



## SEQUENCE LISTING

&lt;110&gt; CPRO-DLO

Agriculture and Agri-food Canada

<120> Use of the BNM3 Transcriptional Activator to Control  
Plant Embryogenesis and Regeneration Processes

&lt;130&gt; E158138

&lt;140&gt;

&lt;141&gt;

&lt;160&gt; 12

&lt;170&gt; PatentIn Ver. 2.1

&lt;210&gt; 1

&lt;211&gt; 2014

&lt;212&gt; DNA

&lt;213&gt; Brassica napus

&lt;400&gt; 1

```

gttcactctct cttctttaag accaaaaacct tttctctctc ctcttcctgc atgaaccccta 60
actaagttct tcttctttta ctttttacca agaactcgtt agatcactct ctgaactcaa 120
tgaataataa ctggttaggc tttctctctc ctcttatga acaaatcac catcgtaagg 180
acgtctactc ttccaccacc acaaccgtcg tagatgtcgc cggagagtac tgttagcgtc 240
cgaccgtcgc ctccgatgag tcttcagcca tccaaacatc gttccttctc cctttgtgtg 300
tcgtcgtcga tgccttcacc agagacaaca atagtcactc ccgagattgg gacatcaatg 360
gttgtgcatg caataacatc cacaacgatg agcaagatgg accaaagctt gagaatttcc 420
ttggccgcac caccacgatt tacaacacca acgaaaacgt tggagatgga agtgggaagtg 480
gctgttatgg aggaggagac ggtggtgtg gctcactagg actttcgatg ataaagacat 540
ggctgagaaa tcaaccctgt gataatgttg ataatacaga aaatggcaat gctgcaaaag 600
gctgtctctc ctcaatgaa ctctctactc cttgtgataa caacaacgac agcaataaca 660
acgtgtgtgc ccaagggaag actattgatg atagcgttga agctacaccg aagaaaaacta 720
ttgagagtgt ttgacagagg acgtctatat accgcggtgt tacaaggcat cgggtgacag 780
gaagatatga ggcacattta tgggataata gttgtaaaag agaaggccaa acgcgcaaaag 840
gaagacaagt ttatttggga ggttatgaca aagaagaaaa agcagctagg gcttatgatt 900
tagccgcact caagtattgg ggaaccacca ctactactaa ctctcccatg agcgaatatg 960
aaaaagaggt agaagagatg aagcacatga caaggcaaga gtatgttgcc tcaactcgca 1020
ggaaaagtat tggtttctct cgtggtgcac cgatttatcg tggagtaaca agacatcacc 1080
aacatggaag atggcaagct aggataggaa gactgcgcgg taacaaagac ctctactttg 1140
gaacttttgg cacacaagaa gaagctgcag aggcatacga cattgcggcc atcaaatcca 1200
gaggattaac cgcagtgact aacttcgaca tgaacagata caacgttaaa gcaatcctcg 1260
aaagccctag tcttctctatt ggtagcgcgc caaaacgtct caaggaggct aaccgtccgg 1320
ttccaagtat gatgatgac agtaataacg tttcagagag tgagaatagt gctagcggtt 1380
ggcnaaacgc tgcggttcag catcatcagg gactagattt gactctattg caccacatc 1440
aaggagagta caatggttat tattacaatg gaggaacctt gtcttcggag agtgctaggg 1500

```

```

cttgtttcaa acaagaggat gatcaacacc atttcttgag caacacgcag agcctcatga 1560
ctaataatcga tcatcaaagt tctgtttcgg atgattcggg tactgtttgt ggaatgttg 1620
ttggttatgg tggttatcaa ggatttgcag ccccggttaa ctgcgatgcc tacgtctgcta 1680
gtgagtttga ttaataacgca agaaaccatt attactttgc tcagcagcag cagaccagc 1740
agtcgccagg tggagatttt cccgcggcaa tgacgaataa tgttggctct aatatgtatt 1800
accatgggga aggtgggtgga gaattgtctc caacatttacc agtttgggaac gacaattaga 1860
aaaaatagtt aaagatcttt agttatatgc gttgtgtgtg gctggtgaac agtgtgatac 1920
tttgattatg ttttttctt tctcttttct tttttcttgg ttaatttctt aagacttatt 1980
tttagtttcc attagtggga taaattttca gact 2014

```

&lt;210&gt; 2

&lt;211&gt; 579

&lt;212&gt; PRT

&lt;213&gt; Brassica napus

&lt;400&gt; 2

```

Met Asn Asn Asn Trp Leu Gly Phe Ser Leu Ser Pro Tyr Glu Gln Asn
  1              5              10              15

```

```

His His Arg Lys Asp Val Tyr Ser Ser Thr Thr Thr Thr Val Val Asp
              20              25              30

```

```

Val Ala Gly Glu Tyr Cys Tyr Asp Pro Thr Ala Ala Ser Asp Glu Ser
      35              40              45

```

```

Ser Ala Ile Gln Thr Ser Phe Pro Ser Pro Phe Gly Val Val Val Asp
      50              55              60

```

```

Ala Phe Thr Arg Asp Asn Asn Ser His Ser Arg Asp Trp Asp Ile Asn
      65              70              75              80

```

```

Gly Cys Ala Cys Asn Asn Ile His Asn Asp Glu Gln Asp Gly Pro Lys
      85              90              95

```

```

Leu Glu Asn Phe Leu Gly Arg Thr Thr Thr Ile Tyr Asn Thr Asn Glu
      100              105              110

```

```

Asn Val Gly Asp Gly Ser Gly Ser Gly Cys Tyr Gly Gly Gly Asp Gly
      115              120              125

```

```

Gly Gly Gly Ser Leu Gly Leu Ser Met Ile Lys Thr Trp Leu Arg Asn
      130              135              140

```

```

Gln Pro Val Asp Asn Val Asp Asn Gln Glu Asn Gly Asn Ala Ala Lys
      145              150              155              160

```

```

Gly Leu Ser Leu Ser Met Asn Ser Ser Thr Ser Cys Asp Asn Asn Asn

```

165	170	175
Asp Ser Asn Asn Asn Val Val Ala Gln Gly Lys Thr Ile Asp Asp Ser		
180	185	190
Val Glu Ala Thr Pro Lys Lys Thr Ile Glu Ser Phe Gly Gln Arg Thr		
195	200	205
Ser Ile Tyr Arg Gly Val Thr Arg His Arg Trp Thr Gly Arg Tyr Glu		
210	215	220
Ala His Leu Trp Asp Asn Ser Cys Lys Arg Glu Gly Gln Thr Arg Lys		
225	230	235
Gly Arg Gln Val Tyr Leu Gly Gly Tyr Asp Lys Glu Glu Lys Ala Ala		
245	250	255
Arg Ala Tyr Asp Leu Ala Ala Leu Lys Tyr Trp Gly Thr Thr Thr Thr		
260	265	270
Thr Asn Phe Pro Met Ser Glu Tyr Glu Lys Glu Val Glu Glu Met Lys		
275	280	285
His Met Thr Arg Gln Glu Tyr Val Ala Ser Leu Arg Arg Lys Ser Ser		
290	295	300
Gly Phe Ser Arg Gly Ala Ser Ile Tyr Arg Gly Val Thr Arg His His		
305	310	315
Gln His Gly Arg Trp Gln Ala Arg Ile Gly Arg Val Ala Gly Asn Lys		
325	330	335
Asp Leu Tyr Leu Gly Thr Phe Gly Thr Gln Glu Glu Ala Ala Glu Ala		
340	345	350
Tyr Asp Ile Ala Ala Ile Lys Phe Arg Gly Leu Thr Ala Val Thr Asn		
355	360	365
Phe Asp Met Asn Arg Tyr Asn Val Lys Ala Ile Leu Glu Ser Pro Ser		
370	375	380
Leu Pro Ile Gly Ser Ala Ala Lys Arg Leu Lys Glu Ala Asn Arg Pro		
385	390	395
Val Pro Ser Met Met Met Ile Ser Asn Asn Val Ser Glu Ser Glu Asn		
405	410	415
Ser Ala Ser Gly Trp Gln Asn Ala Ala Val Gln His His Gln Gly Val		

420

425

430

Asp Leu Ser Leu Leu His Gln His Gln Glu Arg Tyr Asn Gly Tyr Tyr  
435 440 445

Tyr Asn Gly Gly Asn Leu Ser Ser Glu Ser Ala Arg Ala Cys Phe Lys  
450 455 460

Gln Glu Asp Asp Gln His His Phe Leu Ser Asn Thr Gln Ser Leu Met  
465 470 475 480

Thr Asn Ile Asp His Gln Ser Ser Val Ser Asp Asp Ser Val Thr Val  
485 490 495

Cys Gly Asn Val Val Gly Tyr Gly Gly Tyr Gln Gly Phe Ala Ala Pro  
500 505 510

Val Asn Cys Asp Ala Tyr Ala Ala Ser Glu Phe Asp Tyr Asn Ala Arg  
515 520 525

Asn His Tyr Tyr Phe Ala Gln Gln Gln Gln Thr Gln Gln Ser Pro Gly  
530 535 540

Gly Asp Phe Pro Ala Ala Met Thr Asn Asn Val Gly Ser Asn Met Tyr  
545 550 555 560

Tyr His Gly Glu Gly Gly Gly Glu Val Ala Pro Thr Phe Thr Val Trp  
565 570 575

Asn Asp Asn

<210> 3

<211> 2011

<212> DNA

<213> Brassica napus

<400> 3

ttcttctttt accttttacc aagaactcgt tagatcattt tctgaactcg atgaataata 60  
actggttagg cttttctctc tctccttatg aacaaaaatca ccatcgtaag gacgtctgct 120  
cttccaccac cacaaccgcc gtgatgtctg ccggagagta ctgttacgat ccgaccgctg 180  
cctccgatga gtcttcagcc atccaaacat cgtttcccttc tccctttgggt gtcgtctctg 240  
atgctttcac cagagacaac aatagtcact ccgagattg ggacatcaat ggtagtgcac 300  
gtaataacat ccaaatgat gagcaagatg gacccaaact tgagaatttc cttggccgca 360  
ccaccacgat ttacaacacc aacgaaaacg ttggagatat cgatggaagt ggggtgttatg 420  
gaggaggaga cggtggtggt ggctcactag gactttcgtg gataaagaca tggctgagaa 480

```

atcaaccgt ggataatgtt gataatcaag aaaatggcaa tggtgcaaaa ggcctgtccc 540
tctcaatgaa ctcatctact tcttgtgata acaacaacta cagcagtaac aaccttgttg 600
cccaaggga gactattgat gatagcgttg aagctacacc gaagaaaact attgagagtt 660
ttggacagag gacgtctata taccgcggtg ttacaaggca tcggtggaca ggaagatatg 720
aggcacattt atgggataat agttgtaaac gagaaggcca aacgcgcaaa ggaagacaag 780
tttatttggg aggttatgac aaagaagaaa aagcagctag ggcttatgat ttgcccgcac 840
tcaagtattg gggaaccacc actactacta acttccccat gagcgaatat gagaagaga 900
tagaagagat gaagcacatg acaaggcaag agtatgttgc ctccactcgc aggaaaagta 960
gtggtttctc tcgtgttgca tcgatttata gtggagtaac aagacatcac caacatggaa 1020
gatggcaagc taggatagga agagtcgccc gtaacaaaga cctctacttg ggaacttttg 1080
gcacacaaga agaagctgca gaggcatatg acattgcggc catcaaatcc agagagttat 1140
ccgcagtgac taacttcgac atgaacagat acaacgttaa agcaatcctc gaaagcccta 1200
gtcttctcat tggtagcgcc gcaaaacgtc tcaaggaggc taaccgtccg gttccaagta 1260
tgatgatgat cagtaataac gtttcagaga gtgagaataa tgctagcggg tggcaaaacg 1320
ctcgcgttca gcctcatcag ggagtagatt tgagcttatt gcagcaacat caagagaggt 1380
acaatggtta ttattacaat ggaggaaact tgccttcgga gagtgttagg gcttggttca 1440
aacaagagga tgatcaacac catttcttga gcaacacgca gagcctcatg actaatatcg 1500
atcataaag ctctgtttca gatgattcgg ttactgtttg tggaatagtt gttggttatg 1560
gtggtttatca aggatttgca gccccggtta actgcgatgc ctacgtgtgt agtgagttg 1620
actataacgc aagaaccat tattactttg ctacgagca gcagaccag catcgcgacg 1680
gaggagattt tcccgcgcca atgacgaata atgttgctc taatatgata taccatgggg 1740
aagggtggtg agaagttgct ccaacattta cagtttggaa cgacaattag aaataatagt 1800
taaagatctt tagttatatg cgttgtttg tggtgttgaa cagtttgata ctttgattat 1860
gtttttttt ctctttttca ttttgttgt tagttttcta agacttattt tttgtttcca 1920
ttagttggat aaattttcgg acttaagggt cacttctgtt ctgactcttg tctaatacac 1980
aaaagttttc ataaaaaaa aaaaaaaaaa a 2011

```

&lt;210&gt; 4

&lt;211&gt; 579

&lt;212&gt; PRT

&lt;213&gt; Brassica napus

&lt;400&gt; 4

```

Met Asn Asn Asn Trp Leu Gly Phe Ser Leu Ser Pro Tyr Glu Gln Asn
  1             5             10             15

```

```

His His Arg Lys Asp Val Tyr Ser Ser Thr Thr Thr Thr Val Val Asp
      20             25             30

```

```

Val Ala Gly Glu Tyr Cys Tyr Asp Pro Thr Ala Ala Ser Asp Glu Ser
      35             40             45

```

```

Ser Ala Ile Gln Thr Ser Phe Pro Ser Pro Phe Gly Val Val Val Asp
      50             55             60

```

```

Ala Phe Thr Arg Asp Asn Asn Ser His Ser Arg Asp Trp Asp Ile Asn
      65             70             75             80

```

Gly Cys Ala Cys Asn Asn Ile His Asn Asp Glu Gln Asp Gly Pro Lys  
 85 90 95

Leu Glu Asn Phe Leu Gly Arg Thr Thr Thr Ile Tyr Asn Thr Asn Glu  
 100 105 110

Asn Val Gly Asp Gly Ser Gly Ser Gly Cys Tyr Gly Gly Gly Asp Gly  
 115 120 125

Gly Gly Gly Ser Leu Gly Leu Ser Met Ile Lys Thr Trp Leu Arg Asn  
 130 135 140

Gln Pro Val Asp Asn Val Asp Asn Gln Glu Asn Gly Asn Ala Ala Lys  
 145 150 155 160

Gly Leu Ser Leu Ser Met Asn Ser Ser Thr Ser Cys Asp Asn Asn Asn  
 165 170 175

Asp Ser Asn Asn Asn Val Val Ala Gln Gly Lys Thr Ile Asp Asp Ser  
 180 185 190

Val Glu Ala Thr Pro Lys Lys Thr Ile Glu Ser Phe Gly Gln Arg Thr  
 195 200 205

Ser Ile Tyr Arg Gly Val Thr Arg His Arg Trp Thr Gly Arg Tyr Glu  
 210 215 220

Ala His Leu Trp Asp Asn Ser Cys Lys Arg Glu Gly Gln Thr Arg Lys  
 225 230 235 240

Gly Arg Gln Val Tyr Leu Gly Gly Tyr Asp Lys Glu Glu Lys Ala Ala  
 245 250 255

---

Arg Ala Tyr Asp Leu Ala Ala Leu Lys Tyr Trp Gly Thr Thr Thr Thr  
 260 265 270

Thr Asn Phe Pro Met Ser Glu Tyr Glu Lys Glu Val Glu Glu Met Lys  
 275 280 285

His Met Thr Arg Gln Glu Tyr Val Ala Ser Leu Arg Arg Lys Ser Ser  
 290 295 300

Gly Phe Ser Arg Gly Ala Ser Ile Tyr Arg Gly Val Thr Arg His His  
 305 310 315 320

Gln His Gly Arg Trp Gln Ala Arg Ile Gly Arg Val Ala Gly Asn Lys  
 325 330 335

Asp Leu Tyr Leu Gly Thr Phe Gly Thr Gln Glu Glu Ala Ala Glu Ala  
340 345 350

Tyr Asp Ile Ala Ala Ile Lys Phe Arg Gly Leu Thr Ala Val Thr Asn  
355 360 365

Phe Asp Met Asn Arg Tyr Asn Val Lys Ala Ile Leu Glu Ser Pro Ser  
370 375 380

Leu Pro Ile Gly Ser Ala Ala Lys Arg Leu Lys Glu Ala Asn Arg Pro  
385 390 395 400

Val Pro Ser Met Met Met Ile Ser Asn Asn Val Ser Glu Ser Glu Asn  
405 410 415

Ser Ala Ser Gly Trp Gln Asn Ala Ala Val Gln His His Gln Gly Val  
420 425 430

Asp Leu Ser Leu Leu His Gln His Gln Glu Arg Tyr Asn Gly Tyr Tyr  
435 440 445

Tyr Asn Gly Gly Asn Leu Ser Ser Glu Ser Ala Arg Ala Cys Phe Lys  
450 455 460

Gln Glu Asp Asp Gln His His Phe Leu Ser Asn Thr Gln Ser Leu Met  
465 470 475 480

Thr Asn Ile Asp His Gln Ser Ser Val Ser Asp Asp Ser Val Thr Val  
485 490 495

Cys Gly Asn Val Val Gly Tyr Gly Gly Tyr Gln Gly Phe Ala Ala Pro  
500 505 510

Val Asn Cys Asp Ala Tyr Ala Ala Ser Glu Phe Asp Tyr Asn Ala Arg  
515 520 525

Asn His Tyr Tyr Phe Ala Gln Gln Gln Gln Thr Gln Gln Ser Pro Gly  
530 535 540

Gly Asp Phe Pro Ala Ala Met Thr Asn Asn Val Gly Ser Asn Met Tyr  
545 550 555 560

Tyr His Gly Glu Gly Gly Gly Glu Val Ala Pro Thr Phe Thr Val Trp  
565 570 575

Asn Asp Asn

<210> 5  
<211> 4873  
<212> DNA  
<213> Brassica napus

<220>  
<221> intron  
<222> (1846)..(2298)

<220>  
<221> intron  
<222> (2720)..(2952)

<220>  
<221> intron  
<222> (3036)..(3160)

<220>  
<221> intron  
<222> (3170)..(3314)

<220>  
<221> intron  
<222> (3404)..(3553)

<220>  
<221> intron  
<222> (3628)..(3797)

<220>  
<221> intron  
<222> (3849)..(3961)

<220>  
<221> intron  
<222> (4039)..(4148)

<220>  
<221> misc\_feature  
<222> (1620)..(1622)  
<223> start codon

<220>  
<221> misc\_feature  
<222> (4856)..(4858)



&lt;223&gt; stop codon

&lt;400&gt; 5

atctctccac cgattcgtta cccagtgctt gaaaatatga tgactacgaa tcaattaaat 60  
 ggagaagctc cactgctgtt gtaggtggaa gctcaagcaa caaccggaaa cctcgccggt 120  
 atcgggagtt agcatcggtta ttgcgcaaaa ttccgcgcgc agagatgaaa cgattcaaga 180  
 gaaacccotca aataggttag ccatataaaca gtgaattagt atgatttaag agataaaga 240  
 agaagatgag ttcaagaaaa gaaatactca catctattta tactgtttac acaccgcctt 300  
 tcgaactctaa ccaagcattt gaagatgaat cgtggaggag agttaatagg atttaacaca 360  
 aagccattaa ccaaacctgt gcaggtcgag agacgaaccg caaagctcac gcttagccgt 420  
 cgcacgaaga ggagcgatga atttcgtttt ctgcgtgcag tcgtattagg gatagacgga 480  
 gctcattatc gttgggccgg aaacacttct aatctcacag cccatgaaca cactaagaa 540  
 cgaaaccgaa aatgtttgaa gttaaataag acgtgcgggt tgccttagg acacatgtca 600  
 ttacgatag aaatgattta tctacgtgga tcatagggtg ctctctaagg agagagcaaa 660  
 ccttaacttt atataaatag atttgtatca ttctaagag tgtttaagat ttttgcaata 720  
 atattaaaaa aaatacaaaa tttttatgta attagttttg gttacataaa ataacattaa 780  
 ataaaaatfa ttcaaccaat aaaaaatac ggtattttat aattggtcaa aataaaaaat 840  
 aaaaacttaa attcaccta gaattacgag atgtgcactt atttgaaac aaaaataaaa 900  
 tcttttaaca tcaattaaac tgatcggagt ggagtataa tctttacaga gaacatatat 960  
 atatgttttt cttgtaagcg tccatctctt cttagtcagt tagttcaaat accagcttga 1020  
 gtaaaacctt gaattttgta atttgttgta aaatattcga agcgactact cactagtggt 1080  
 aagcaaaacg ccaaacgcaa tgcgtcgctc ggtcataggg tcaacacatc acatgtgact 1140  
 agcattatgg gtcttaattc aacagcgagt gattttggga tttattatta gttctcgtgt 1200  
 tactctact ttacacaaa gtccataacc ttatttacac atgaagagag ttttgcaagg 1260  
 gcttttgact gattaattat aatgtattaa accaaactag aattaagaga ttaggcattg 1320  
 aattacatta ccaccaccac ccaccattca aaccgacca tacatctcca cagttttcaa 1380  
 gtaaaacaaa ttttttttgt tgttccctcg gaatttaaat aaatattcgt ttataataa 1440  
 gcgcgatgata tgacgcctcg gaagaaatga aacatttatat ctttgacttt tcttctccta 1500  
 gttcatctct cttctttaag accaaaacct ttttctctc ctcttcatgc atgaacccta 1560  
 actaagttct tctcttttta ccttttacca agaactcggt agatcactct ctgaactcaa 1620  
 tgaataataa ctggttaggc ttttctctct ctcttcatga acaaaatcac catcgtaagg 1680  
 acgtctactc ttccaccacc acaaccgtcg tagatgtcgc cggagagtac tgttacgac 1740  
 gcagccgtcg ctccgatgag tcttcagcca tccaaacatc gtttctctct ccttttggtg 1800  
 tctgtctgga tcttttccac agagacacaa atgtctctc cggaggttat tgtttbaga 1860  
 ctacttgatt ttttttgatt tgtttatttg tttagtttcc tcttcttcca atgcgtagaa 1920  
 caaagaccaa tacacacgca cgcatactag cctatttttt tcttgggtct cctatgaga 1980  
 tttcatttat ttgagaata tcaatgtgtg gggtttgatg tttgttgca tatagtaata 2040  
 ctaaaacata tgccagttat acatagattt tttttaaaga tatacatgga tatgaaatga 2100  
 aatttgacat ttctctcttt attcaatata ataatatgat cacatacatg tgaactttt 2160  
 gatttgataa tttgttctt acagtgaag gagagaataa ccaaatcccc atttgataa 2220  
 tatagatctg tgatgaaaaa taaatttaac aaattatgat aatataggcc attaatcttt 2280  
 gttttttttt ctttatagat tgggacatca atggttgtgc atgcaataac atcccaatcg 2340  
 atgacgaaga tggaccaaag cttgagaatt tcttggcccg caccaccacg atttacaaca 2400  
 ccaacgaaaa cggtggagat ggaagtggaa gtggttgta tggaggagga gacggtggtg 2460  
 gtggtctact aggaactttcg atgataaaga catggtcgag aaatcaaccc ttggataaag 2520  
 ttgataatca agaaaatggc aatgctgcaa aaggcctgtc cctctcaatg aactcatcta 2580  
 cttcttttga taacaacac gacagcaata acaacgttgt tgcccaaggg aagactattg 2640  
 atgatagctg tgaagctaca ccgaagaaaa ctatgagag ttttgagacg aggcgtcta 2700

```

tataccgcgg tgttacaagg tgccttcat ttatttaatt aaaatgtgta aaatgtcgct 2760
tgaattgtta tcttcttggg aaagtctggg acattgactc aatggctctg ttgcgagagt 2820
gctaccggaat ggtccttgat atatagatc aaagagagat attgttatta tgggcttata 2880
tagaataata catatatata tatatatata tggtagctgt tgatgacatg tatgttcgta 2940
ttaaattgata aggcattcggg ggacaggaag atatgagcca ctttatggg ataattgttg 3000
taaaagagaa ggccaaacgc gcaaaggaag acaaggtata tatatttca ttgataattt 3060
gatcatattt tcatacacga ttactttcca aactaatata ggttttccga tcaattgtcca 3120
tggtttttatc aaaattttgca cctgggtggtt gtcttctcag ttattttggg taagtaattt 3180
attataaatt ggacgaagct gtgatgggta aatctaaatt atataatcaa atttgtttat 3240
tttttggta tacattccatt atataatcaa aatagcgata cgatctacat caaattgttg 3300
tctatatcat gcagagggtt atgacaaaga agaaaaagca gctaggcgct atgatttagc 3360
cgactccaag tattggggaa ccaccactac tactaaactc ccgtaagtc aatcaatgtt 3420
gtacaagatt tcataactta gaaccaattt tattcttttt ttataagatg ctattatcct 3480
attattaatt gccattgttta tatcgttaca ttattacaa taaaagtagc ttttggtttg 3540
atataatag tagatgagcg aatatgaaa agaggtagaa gagatgaagc acatgacaag 3600
gcaagagatg gttgcctcac tgcgcaggta tataatggaa cttctgatat tattgcatat 3660
ggcactctatt attatacatg tatattagta tttatatat agaaccatc acgctcacgt 3720
ttatttttaa aaatatgtcc gtattcacgt cagattatca gcatacacct atataataa 3780
gacataaaaa tatgcaggaa aagtagtggt ttctctcggt gtgcacgat ttatcgtgga 3840
gtaacaaggt attcatcacg agagaacgaa tcctattttg ttactgtacat atatatataa 3900
aaatataatt ataagatata acattttata ttatgaatat ttcttctaatt ggggtccaaa 3960
gacatacaca acatggaaga tggcaagcta ggtatggaag agtcgccggt aacaagacc 4020
tctacttggg aacttttgggt acgttttagtc ttctcttact aaacttcaca atcaaateta 4080
taacaaaaga tatcaactaa aaactacaac atatatctaa gtaagctgta catattatt 4140
atatgaagcg acacaagaag aagctgcaga ggcatacgac attgcggcca tcaaatccag 4200
aggattaacc gcagtgacta acttcgacat gaacagatac aacgttaaag caatcctcga 4260
aagccctagt cttcctattg gtacgcgcgc aaaaacgtctc aaggaggcta accgtccggt 4320
tccaagtatg atgatgatca gtaataacgt ttcagagagt gagaatagtg ctacggttg 4380
gcaaaacgct gcggttcagc atcatcagg agtagatttg agcttattgc accaatcaca 4440
agagaggatc aatggttatt attacaatgg aggaacttg tcttcggaga gtgctagggc 4500
ttgtttcaaa caagaggatg atcaacacca ttctctgagc aacacgcaga gcctcatgac 4560
taatatcgat catcaaaagt ctgtttcgga tgattcggtt actgtttgtg gaaatgttgt 4620
tggttatggt ggttatcaag gatttcgac ccggttaac tgcgatgcct acgctgctag 4680
tgagttgat tataacgaa gaacattta ttactttgtt agagagagc agacccagca 4740
gtcgcagggt ggagattttc ccgcggcaat gacgaataat gttggctcta atatgtatta 4800
ccatggggaa ggtggtggag aagttgctcc aacatttaca gtttggaacg acaattagaa 4860
aaaatagtta aag 4873

```

&lt;210&gt; 6

&lt;211&gt; 30

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:primer

&lt;400&gt; 6

gaggcagcgg tcggatcgta acagtactct

30

&lt;210&gt; 7

&lt;211&gt; 30

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:primer

&lt;400&gt; 7

cataaggaga gagagaaaag cctaaccagt

30

&lt;210&gt; 8

&lt;211&gt; 19

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:primer

&lt;400&gt; 8

accaagaact cgtagatc

19

&lt;210&gt; 9

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:primer

&lt;400&gt; 9

aacgcatata actaaagatc

20

&lt;210&gt; 10

&lt;211&gt; 26

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:primer

&lt;400&gt; 10

ccatggatcc agagacgaag cgaaac

26

<210> 11

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 11

actccatgga taataactgg ttaggc

26

<210> 12

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:primer

<400> 12

aaattctcaa gctttgtcc atcttg

26

- 61 -

EPO - DG 1

02.06.1999

(58)

## CLAIMS

1. An isolated DNA molecule comprising a nucleotide sequence that hybridizes to SEQ ID NO:5 under stringent conditions.
2. The isolated DNA molecule of claim 1 wherein said isolated DNA molecule comprises at least 23 contiguous nucleotides of SEQ ID NO:5.
3. The isolated DNA molecule of claim 1 wherein said isolated DNA molecule comprises a nucleotide sequence that is at least 70% homologous with the nucleotide sequence defined by SEQ ID NO:5.
4. An isolated DNA molecule comprising a nucleic acid sequence encoding a protein, wherein said protein when present at a sufficient level within a plant cell renders said cell embryogenic, increases the regenerative capacity of said plant cell, or both renders said plant cell embryogenic and increases the regenerative capacity of said plant cell, said isolated DNA molecule having at least 70% homology within a nucleotide sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3 and SEQ ID NO:5.
5. The isolated DNA molecule of claim 4 comprising a nucleotide sequence that hybridizes to nucleotides 1620-4873 of SEQ ID NO:5 under stringent conditions.
6. The isolated DNA molecule of claim 4 comprising a nucleotide sequence that hybridizes to the nucleotide sequence of SEQ ID NO:1 under stringent conditions.

- 62 -

7. The isolated DNA molecule of claim 4 comprising a nucleotide sequence that hybridizes to the nucleotide sequence of SEQ ID NO:3 under stringent conditions.
8. The isolated DNA molecule of claim 6, wherein said DNA encodes a protein as defined by SEQ ID NO:2.
9. The isolated DNA molecule of claim 7, wherein said DNA encodes a protein as defined by SEQ ID NO:4.
10. A vector comprising the isolated DNA molecule as claimed in any one of claims 1 to 9, wherein said isolated DNA molecule is under control of a regulatory element that directs expression of said DNA in a plant cell.
11. The vector of claim 10, wherein said regulatory element is a constitutive regulatory element
12. The vector of claim 10, wherein said regulatory element is an inducible regulatory element.
13. The vector of claim 10, wherein said regulatory element is a tissue specific regulatory element
14. The vector of claim 10, wherein said regulatory element is an developmentally active regulatory element.
15. A transformed plant cell comprising the vector of any one of claims 10 to 14.
16. A transformed plant comprising the vector of any one of claims 10 to 14.
17. A seed obtained from the transformed plant of claim 16.

- 63 -

18. An isolated protein encoded by the isolated DNA molecule as claimed in any one of claims 4 to 9.
19. A method of producing asexually derived embryos comprising:
- i) transforming a plant cell with the vector of any one of claims 10 to 14;
  - ii) growing said plant cell to produce transformed tissue;
  - iii) selecting said transformed tissue for occurrence of said isolated DNA molecule; and
  - iv) assaying said transformed plant for asexual embryo production.
20. The method of claim 19 wherein the step of assaying involves assaying for adventitious embryony.
21. The method of claim 19, wherein the step of assaying involves assaying for somatic embryos.
22. The method of claim 19, wherein the step of assaying involves assaying for gametophytic embryos.
23. The method of claim 19, wherein the step of assaying involves assaying for haploid parthenogenesis of the embryo sac.
- 
24. The method of claim 19, wherein the step of assaying involves assaying for diplospory.
25. A method of modifying the regenerative capacity of a plant comprising
- i) transforming a plant cell with the vector of any one of claims 10 to 14;
  - ii) growing said transformed plant cell to produce transformed tissue; and

- 64 -

- iii) assaying said transformed plant tissue for enhanced regeneration as compared to wild-type tissue.
26. The method of claim 25, wherein the step of growing said transformed plant cell, the step of assaying said transformed plant tissue, or both the step of growing said transformed plant cell and the step of assaying said transformed plant tissue are carried out in the absence of a growth regulator.
27. A method of selecting a transformed plant comprising:
- i) transforming a normally non-regenerative plant with a vector of any one of claims 10 to 14; and
  - ii) determining whether said transformed plant is able to regenerate under conditions in which said normally non-regenerative plant does not regenerate.
28. The isolated DNA molecule of claim 1 comprising a DNA sequence that hybridizes to nucleotides 1-1619 of SEQ ID NO:5 under stringent conditions.
29. The isolated DNA molecule of claim 1 wherein said isolated molecule comprises at least 22 contiguous nucleotides within nucleotides 1-1619 of SEQ ID NO:5.
- 
30. A vector comprising the isolated DNA molecule of either claim 28 or 29 operably associated with a gene of interest, wherein said isolated DNA molecule directs the expression of said gene of interest within a plant cell.
31. The vector as defined by claim 30, wherein said gene of interest is heterologous with respect to the isolated DNA molecule.
32. The vector as defined by claim 31, wherein said gene of interest is selected from the group consisting of a pharmaceutically active protein, antibody,



- 65 -

industrial enzyme, protein supplement, nutraceutical, storage protein, animal feed and animal feed supplement.

33. A transformed plant cell comprising the vector of either claim 30, 31 or 32.
34. A transformed plant comprising the vector of either claim 30, 31 or 32.
35. A seed obtained from the transformed plant of claim 34.
36. A method for directing the expression of a gene of interest within a developing embryo of a plant comprising transforming said plant with the vector as defined by either claim 30, 31 or 32.
37. A use of a nucleotide sequence as defined in any one of claims 4, 5, 6 or 7 as a selectable marker.
38. A method of producing asexually derived embryos comprising:
  - i) transiently transforming a plant cell with the vector of any one of claims 10 to 14, or introducing into said plant cell the protein of claim 18, to produce a modified plant cell;
  - ii) ~~growing said modified plant cell to produce tissue; and~~
  - iii) assaying said tissue for asexual embryo formation.
39. The method of claim 38 wherein the step of assaying involves assaying for adventitious embryony.
40. The method of claim 38, wherein the step of assaying involves assaying for somatic embryos.
41. The method of claim 38, wherein the step of assaying involves assaying for gametophytic embryos.

- 66 -

42. The method of claim 38, wherein the step of assaying involves assaying for haploid parthenogenesis of the embryo sac.
43. The method of claim 38, wherein the step of assaying involves assaying for diplospory.
44. A method of modifying the regenerative capacity of a plant comprising
- i) transiently transforming a plant cell with the vector of any one of claims 10 to 14 or introducing into said plant cell the protein of claim 18, to produce a modified plant cell;
  - ii) growing said modified plant cell to produce tissue; and
  - iii) assaying said tissue for enhanced regeneration as compared to wild-type tissue.
45. The method of claim 44, wherein the step of growing said modified plant cell, the step of assaying said tissue, or both the step of growing said modified plant cell and the step of assaying said tissue are carried out in the absence of a growth regulator.
- 
46. A method of producing an apomictic plant comprising:
- i) transforming a plant with the vector of any one of claims 10 to 14, to produce a transformed plant;
  - ii) selecting said transformed plant for occurrence of said isolated DNA molecule; and
  - iii) assaying said transformed plant for asexual embryo production.
47. The method of claim 46 wherein the step of assaying involves assaying for adventitious embryony.

- 67 -


48. The method of claim 46, wherein the step of assaying involves assaying for somatic embryos.
49. The method of claim 46, wherein the step of assaying involves assaying for gametophytic embryos.
50. The method of claim 46, wherein the step of assaying involves assaying for parthenogenesis of the embryo sac.
51. A method of modifying the regenerative capacity of a plant comprising
- i) transiently transforming a plant cell with the vector of any one of claims 10 to 14, or introducing into said plant cell the protein of claim 18;
  - ii) growing said plant cell to form tissue; and
  - iii) assaying said tissue for enhanced regeneration as compared to wild-type tissue.
52. The method of claim 51, wherein the step of growing said plant cell, the step of assaying said tissue, or both the step of growing said plant cell and the step of assaying said tissue are carried out in the absence of a growth regulator.
- 
53. A method of selecting a modified plant comprising;
- i) transiently transforming a normally non-regenerative plant with a vector of any one of claims 10 to 14, or introducing into said normally non-regenerative plant the protein of claim 18, to produce said modified plant; and
  - ii) determining whether said modified plant is able to regenerate under conditions in which said normally non-regenerative plant does not germinate.

- 68 -

54. An isolated DNA molecule comprising a sequence encoding a protein consisting of two AP2 DNA binding domains, which when said protein is expressed at a sufficient level in a plant cell, renders said cell embryogenic, or increase the regenerative capacity of said plant cell, or both renders said cell embryogenic and increase the regenerative capacity of said plant cell.
55. A method of producing a protein of interest comprising
- i) transforming a plant with at least one vector, said at least one vector selected from any one of claims 10 to 14 to produce a transformed plant;
  - ii) selecting said transformed plant for occurrence of said isolated DNA molecule; and
  - iv) growing said transformed plant in order to produce said protein of interest, wherein expression of said protein of interest is induced by the expression product of said isolated DNA.
56. The method of claim 55, wherein said transformed plant is transformed with a second vector comprising a nucleotide sequence encoding said protein of interest under the control of a regulatory element, said regulatory element induced by the expression product of said isolated DNA..
- 
57. The method of claim 55, wherein said protein of interest is a native protein.
58. The method of any one of claims 55 or 56, wherein said protein of interest is selected from the group consisting of a pharmaceutically active protein, antibody, industrial enzyme, protein supplement, nutraceutical, storage protein, an enzyme involved in oil biosynthesis, animal feed, and animal feed supplement.

- 69 -

59. The isolated DNA molecule of claim of any one of claims 4 to 7, wherein said isolated DNA molecule encodes a protein that is at least 70% homologous with the amino acid defined by SEQ ID NO:2.
60. The isolated DNA molecule of claim of any one of claims 4 to 7, wherein said isolated DNA molecule encodes a protein that is at least 70% homologous with the amino acid defined by SEQ ID NO:4.
61. The isolated protein of claim 18, wherein said protein comprises from about 30 to about 541 amino acids of the sequence disclosed in SEQ ID NO:2
62. The isolated protein of claim 18, wherein said protein comprises from about from about 30 to about 561 amino acids of the sequence disclosed in SEQ ID NO: 4.
-




E DG 1

02.06.1999

## ABSTRACT OF THE DISCLOSURE

(58)

The present invention provides for a gene obtained during the induction of microspore embryogenesis. The protein encoded by this gene renders plant cells embryogenic, and increases the regenerative capacity of the plant cell. Also disclosed is the regulatory region of this gene and its use for directing the expression of a gene of interest within a suitable host cell.





1/9

Figure 1




Figure 2

2/9

BNM3A	GTTCATCTCTCTCTTTAAGACCAAAACCTTTTCTCTCTCTTTCATGCATGAACCCCTA	60
BNM3B	-----	
BNM3A	ACTAAGTTCTCTCTCTTTTACCTTTTACCAAGAACTCGTTAGATCACTCTCTGAACCTCAA	120
BNM3B	-----TTCTCTCTTTTACCTTTTACCAAGAACTCGTTAGATCAATTTCTGAACCTCGA	51
BNM3A	TGAATAATAACTGGTTAGGCTTTTCTCTCTCTCTTATGAACAAAATCACCATCGTAAGG	180
BNM3B	-----	111
BNM3A	ACGTCTACTCTTCCACCACCACAACCGTCGTAGATGTCGCCGAGAGTACTGTTACGATC	240
BNM3B	ACGTCTGCTCTTCCACCACCACAACCGCGTAGATGTCGCCGAGAGTACTGTTACGATC	171
BNM3A	CGACCGCTGCTCCGATGAGTCTTCAGCCATCCAAACATCGTTTCCTCTCTCCCTTTGGTG	300
BNM3B	CGACCGCTGCTCCGATGAGTCTTCAGCCATCCAAACATCGTTTCCTCTCTCCCTTTGGTG	231
BNM3A	TCGTCTGTCGATCTTCTTCCACGAGACAAANTAGTCACTCCCGAGATTGGGACATCAATG	360
BNM3B	TCGTCTGTCGATCTTCTTCCACGAGACAAANTAGTCACTCCCGAGATTGGGACATCAATG	291
BNM3A	GTGTGTCATGCAATAACATCCACAACGATGAGCAAGATGGACCAAAAGCTTGAGAATTTC	420
BNM3B	GTATGTCATGTAATAACATCCACAATGATGAGCAAGATGGACCAAACTTGAGAATTTC	351
BNM3A	TTGGCCGACCAACACGATTACAAACCAACGAAACGTTGGAGATGGAAGTGGAAAGTG	480
BNM3B	TTGGCCGACCAACACGATTACAAACCAACGAAACGTTGGAGATGTCGATGGAAGTG	411
BNM3A	CGCTGTTATGGAGGAGGAGACGCTGGTGGTGGCTCACTAGGACTTTCGATGATAAGACAT	540
BNM3B	GGTGTATGGAGGAGGAGACGCTGGTGGTGGCTCACTAGGACTTTCGATGATAAGACAT	471
BNM3A	GGCTGAGAAATCAACCCGTGGATATGTTGATATCAAGAAATGGCAATGCTGCAAAAG	600
BNM3B	GGCTGAGAAATCAACCCGTGGATATGTTGATATCAAGAAATGGCAATGCTGCAAAAG	531
BNM3A	GCCTGTCCCTCTCAATGAACCTCATCTACTTCTGTGATAACAACACGACAGCAATAACA	660
BNM3B	GCCTGTCCCTCTCAATGAACCTCATCTACTTCTGTGATAACAACACGACAGTAACA	591
BNM3A	ACCTTGTGTGCCCAAGGGAAGACTATTGATGATAGCGTTGAAGCTACACCGAAGAAACTA	720
BNM3B	ACCTTGTGTGCCCAAGGGAAGACTATTGATGATAGCGTTGAAGCTACACCGAAGAAACTA	651
BNM3A	TTGAGAGTTTTCACACAGGAGACCTATATACCGCGGTGTACBGGCATCTCGTGACAG	780
BNM3B	TTGAGAGTTTTCACACAGGAGACGCTATATACCGCGGTGTACAGGCACTCGTGACAG	711
BNM3A	GAGATATGAGGCACATTTATGGGATATAGTTGTAAAAGAGAAGGCCAAACGCGCAAAAG	840
BNM3B	GAGATATGAGGCACATTTATGGGATATAGTTGTAAACGAGAAGGCCAAACGCGCAAAAG	771
BNM3A	GAGACAAGTTTATTTGGGAGGTTATGACAAAAGAAAAGACAGCTAGGGCTATGATT	900
BNM3B	GAGACAAGTTTATTTGGGAGGTTATGACAAAAGAAAAGACAGCTAGGGCTATGATT	831
BNM3A	TAGCCGCACTCAAGTATTGGGGAACCACTACTACTACTTCCCATGAGCGAATATG	960
BNM3B	TAGCCGCACTCAAGTATTGGGGAACCACTACTACTACTTCCCATGAGCGAATATG	891
BNM3A	AAAAGAGGTAGAAGAGATGAAGCACATGACAAGGCAAGATGATGTGCTCTACTCGCA	1020
BNM3B	AGAAAGAGTAGAAGAGATGAAGCACATGACAAGGCAAGATGATGTGCTCTACTTCGCA	951
BNM3A	GGAAAAAGTACTGGTTCTCTCTGTTGGTGCATCGATTATCGTGGAGTAACAGACATCACC	1080
BNM3B	GGAAAAAGTACTGGTTCTCTCTGTTGGTGCATCGATTATCGTGGAGTAACAGACATCACC	1011

3/9

BNM3A	AACATGGAAGATGGCAAGCTAGGATAGGAAGAGTCCCGGTAAACAAAGACCTCTACTTGG	1140
BNM3B	AACATGGAAGATGGCAAGCTAGGATAGGAAGAGTCCCGGTAAACAAAGACCTCTACTTGG	1071
*****		
BNM3A	GAACTTTTGGCACAACAAGAAGCTGCAGAGGCATACGACATTGCGGCCATCAAAATTC	1200
BNM3B	GAACTTTTGGCACAACAAGAAGCTGCAGAGGCATACGACATTGCGGCCATCAAAATTC	1131
*****		
BNM3A	GAGGATTAAACCGCAGTGACTAACTTCGACATGAACAGATACAACGTTAAAGCAATCCCTCG	1260
BNM3B	GAGGATTAAACCGCAGTGACTAACTTCGACATGAACAGATACAACGTTAAAGCAATCCCTCG	1191
*****		
BNM3A	AAAGCCCTAGTCTTCTTATTGGTAGCGCCGCAAAACGCTCAAGGAAGCTAACCGTCCGG	1320
BNM3B	AAAGCCCTAGTCTTCTTATTGGTAGCGCCGCAAAACGCTCAAGGAAGCTAACCGTCCGG	1251
*****		
BNM3A	TTCCAAGTATGATGATGATCAGTAATAACGTTTCAGAGAGTGAGAATAGTGTAGCGGTT	1380
BNM3B	TTCCAAGTATGATGATGATCAGTAATAACGTTTCAGAGAGTGAGAATAGTGTAGCGGTT	1311
*****		
BNM3A	GGCAAAACGCTCGCGTTTTCAGCATCATCAGGGAGTAGATTGAGCTTATTGCAACCAATC	1440
BNM3B	GGCAAAACGCTCGCGTTTTCAGCATCATCAGGGAGTAGATTGAGCTTATTGCAACCAATC	1371
*****		
BNM3A	AAGAGAGGTACAACTGTTTATTATTCAAATGAGGAAACCTTCTCTCGAGAGTCTAGGG	1500
BNM3B	AAGAGAGGTACAACTGTTTATTATTCAAATGAGGAAACCTTCTCTCGAGAGTCTAGGG	1431
*****		
BNM3A	CTTGTTCAAACAAGAGGATGATCAACACCATTTCTTGAGCAACACGAGAGCCTCATGA	1560
BNM3B	CTTGTTCAAACAAGAGGATGATCAACACCATTTCTTGAGCAACACGAGAGCCTCATGA	1491
*****		
BNM3A	CTAATATCGATCATCAAAGTTCTGTTTCGGATGATTCGGTTACTGTTTGTGGAAATGTTG	1620
BNM3B	CTAATATCGATCATCAAAGTTCTGTTTCGGATGATTCGGTTACTGTTTGTGGAAATGTTG	1551
*****		
BNM3A	TTGGTTATGGTGTATTCAAGGATTTTCAGCCCGGTTAACTGCGATGCTTACCGCTCTA	1680
BNM3B	TTGGTTATGGTGTATTCAAGGATTTTCAGCCCGGTTAACTGCGATGCTTACCGCTCTA	1611
*****		
BNM3A	GTGAGTTTGATTATAACGCAAGAAACCATTTACTTTGCTCAGCAGCAGCAGACCCAGC	1740
BNM3B	GTGAGTTTGATTATAACGCAAGAAACCATTTACTTTGCTCAGCAGCAGCAGACCCAGC	1671
*****		
BNM3A	AGTCGCCAGGTGGAGATTTTCCCGCGCAATGACGAATAAGTTGGCTCTAATATGTATT	1800
BNM3B	AGTCGCCAGGAGGAGATTTTCCCGCGCAATGACGAATAAGTTGGCTCTAATATGTATT	1731
*****		
BNM3A	ACCATGGGGAAGTGGTGGGAGAAGTTGCTCCAACATTTACAGTTTGAACGACAAATAGA	1860
BNM3B	ACCATGGGGAAGTGGTGGGAGAAGTTGCTCCAACATTTACAGTTTGAACGACAAATAGA	1791
*****		
BNM3A	AAAAATAGTTAAAGATCTTTAGTTATATGCGTTGTTGTGCTGGTGAACAGTGTGATAC	1920
BNM3B	AAAAATAGTTAAAGATCTTTAGTTATATGCGTTGTTGTGCTGGTGAACAGTGTGATAC	1851
*****		
BNM3A	TTTGATTATGTTTTTTTCTTTCTCTTTTCTTTTCTTTGGTTAAATTTCTTAAGACTTATT	1980
BNM3B	TTTGATTATGTTTTTTTCTTTCTCTTTTCTTTTCTTTGGTTAAATTTCTTAAGACTTATT	1909
*****		
BNM3A	TTTAGTTTCCATTAGTTGGATAAAATTTTCAGACT-----	2014
BNM3B	TTTAGTTTCCATTAGTTGGATAAAATTTTCAGACT-----	1969
*****		
BNM3A	-----	
BNM3B	GTCTAATACGAAAGTTTTTCAT	1992

Figure 3

4/9

```

BNM3A MNNNWLGFSLSPYEQNHHRKDVYSSTTTTVDVAGEYCYDPTAASDESSAIQTSFSPFPG 60
BNM3B MNNNWLGFSLSPYEQNHHRKDVCSSTTTTAVDVAGEYCYDPTAASDESSAIQTSFSPFPG
*****

BNM3A VVVDAPTRDNNSHSRDWDINGACNNIHNDQDQPKLENFLGRITTIYNTNENVGDGSGS 120
BNM3B VVVDAPTRDNNSHSRDWDINGACNNIHNDQDQPKLENFLGRITTIYNTNENVGDIDGS
*****

BNM3A GCYGGGDDGGGSLGLSMIKTWLRNPVDNVNDQENGNAAGLSLMMNSSTSCDNNNDSNN 180
BNM3B GCYGGGDDGGGSLGLSMIKTWLRNPVDNVNDQENGNAAGLSLMMNSSTSCDNNNYSNN
*****

repeat 1
BNM3A NVVAQGKTI DDSVEATPKKTI ESFGORTSIYRGVTRHRWTGRYEAHLWDMNSCKREGQTRK 240
BNM3B NLVAQGKTI DDSVEATPKKTI ESFGORTSIYRGVTRHRWTGRYEAHLWDMNSCKREGQTRK
*****

BNM3A GRQVYLGSDYDKEEKAARAYDLAALKYWGTTTTTTFMFSEYEKEVEEMKHMTRQEVVASLR 300
BNM3B GRQVYLGSDYDKEEKAARAYDLAALKYWGTTTTTTFMFSEYEKEVEEMKHMTRQEVVASLR
*****

repeat 2
BNM3A RKSSGFSRGASIRGVTRRHQHGHRWQARIQGVAGNKL DLYLQTFGTQEEAAEAYDLAAIKF 360
BNM3B RKSSGFSRGASIRGVTRRHQHGHRWQARIQGVAGNKL DLYLQTFGTQEEAAEAYDLAAIKF
*****

BNM3A RGLTAVTNFDMRYNVKAILSPSLPIGSAAKRLKEANRPVPSMMMI SNNVSESENSASG 420
BNM3B RGLTAVTNFDMRYNVKAILSPSLPIGSAAKRLKEANRPVPSMMMI SNNVSESENNASG
*****

BNM3A WQNAAVQHHQGVDSL LLLHQHQRERYNGYYNCGNLSSESARACFKQEDDQHFLSNTQSLM 480
BNM3B WQNAAVQHHQGVDSL LLLHQHQRERYNGYYNCGNLSSESARACFKQEDDQHFLSNTQSLM
*****


BNM3A TNIDHQSSVSDDSVTVCGNVVGYGGYQGFAAPVNC DAYAASEFDYNARNHYFAQQQQQTQ 540
BNM3B TNIDHQSSVSDDSVTVCGNVVGYGGYQGFAAPVNC DAYAASEFDYNARNHYFAQQQQQTQ
*****

BNM3A QSPGGDFPAAMTNNVGSNNMYHGEGGGEVAPTFTVWNDN 579
BNM3B HSPGGDFPAAMTNNVGSNNMYHGEGGGEVAPTFTVWNDN
*****

```

5/9

Figure 4



6/9

## Figure 5

## AP2 DOMAIN REPEAT 1

BNM3 TSIYRGVTRHRWGRYEHLWDNSKREGQTRKGQVYLGVDKEEKAARAYDLAALKYWGTTTTNFPWSEYEKEV  
 ANT T\*Q\*\*\*\*\*P\*K\*HS\*\*\*\*\*M\*\*\*\*\*PS\*H\*\*\*SAEN\*Q\*I I  
 ZM T\*\*\*\*\*E\*\*R\*\*\*S\*\*\*\*\*F\*\*P\*\*\*\*\*QV\*N\*\*\*\*L  
 GL15 S\*Q\*\*\*\*\*FY\*R\*\*\*W\*S\*I\*---C-----\*K\*\*\*\*\*P\*TAQA\*\*\*\*\*Q\*I\*FR\*LNADI\*\*TLDD\*KD\*M  
 AP2 S\*Q\*\*\*\*\*FY\*R\*\*\*W\*S\*I\*---C-----\*K\*\*\*\*\*P\*TAHA\*\*\*\*\*R\*I\*FR\*VEADI\*\*NIDD\*DDDL

## LINKER

BNM3 EEMKHMTRQEVYASLRKSSGFSRC  
 ANT \*D\*N\*\*\*\*\*H\*\*\*\*\*  
 ZM \*\*\*\*\*FI\*\*\*\*\*  
 GL15 KK\*\*DLSKEF\*LV\*\*QGA\*\*V\*\*  
 AP2 KQ\*TNLKEE\*HV\*\*Q\*\*T\*\*P\*\*

## AP2 DOMAIN REPEAT 2

BNM3 ASIYRGVTRHHQHGRWQARIGRVAGNKDLYLGTGTQEEAAEAYDIAAIKFRGLAVTNFDMNRVNVKA  
 ANT \*\*\*\*\*V\*\*\*\*\*TN\*\*\*\*\*IT\*\*D\*DR  
 ZM \*\*\*\*\*S\*\*\*\*\*N\*\*\*\*\*L\*\*S\*\*DES  
 GL15 S\*RF\*\*\*\*\*Q\*KC\*K\*E\*\*\*\*\*OLM\*\*K\*YV\*\*LYD\*ET\*\*Q\*\*K\*\*\*CV\*K\*\*\*\*\*AQSD\*DKEL  
 AP2 S\*K\*\*\*\*\*L\*KC\*\*E\*\*M\*QEL\*\*K\*YV\*\*L\*D\*EV\*\*R\*\*K\*\*\*CN\*KD\*\*\*\*\*FSI\*DEEL

7/9

Figure 6

A

+ embryo  
pollen 0 d  
pollen 4 d  
- embryo  
globular  
heart  
torpedo  
21d cot  
28d cot  
42d cot

B



7d 14 d 18 d 21 d 28 d 35 d 42 d

C

leaf  
root  
flower  
small bud  
large bud  
anther  
pistil  
silique

Figure 7

8/9





9/9

Figure 8

